

# Advil Caplets\* (Film Coated)

## Ibuprofen Tablets, 200 mg

This is the most current labeling information, and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

\*capsule-shaped tablets

<b>Drug Facts</b>	
<b>Active ingredient (in each caplet)</b>	<b>Purpose</b>
Ibuprofen 200 mg (provided as ibuprofen sodium 256 mg) (NSAID)* ..... *nonsteroidal anti-inflammatory drug	Pain reliever/Fever reducer
<b>Uses</b>	
<ul style="list-style-type: none"><li>■ temporarily relieves minor aches and pains due to:<ul style="list-style-type: none"><li>■ headache</li><li>■ toothache</li><li>■ backache</li><li>■ menstrual cramps</li><li>■ the common cold</li><li>■ muscular aches</li><li>■ minor pain of arthritis</li></ul></li><li>■ temporarily reduces fever</li></ul>	
<b>Warnings</b>	
<p><b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"><li>■ hives</li><li>■ facial swelling</li><li>■ asthma (wheezing)</li><li>■ shock</li><li>■ skin reddening</li><li>■ rash</li><li>■ blisters</li></ul> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"><li>■ are age 60 or older</li><li>■ have had stomach ulcers or bleeding problems</li><li>■ take a blood thinning (anticoagulant) or steroid drug</li><li>■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]</li><li>■ have 3 or more alcoholic drinks every day while using this product</li><li>■ take more or for a longer time than directed</li></ul> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p>	
<b>Do not use</b>	
<ul style="list-style-type: none"><li>■ if you have ever had an allergic reaction to any other pain reliever/fever reducer</li><li>■ right before or after heart surgery</li></ul>	

# Advil Caplets\* (Film Coated)

## Ibuprofen Tablets, 200 mg

This is the most current labeling information, and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

\*capsule-shaped tablets

### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### When using this product

- take with food or milk if stomach upset occurs

### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

# Advil Caplets\* (Film Coated)

## Ibuprofen Tablets, 200 mg

This is the most current labeling information, and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

\*capsule-shaped tablets

### ***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

### ***Other information***

- each caplet contains: **sodium 22 mg**
- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

### ***Inactive ingredients***

acesulfame potassium, caramel color, carnauba wax, colloidal silicon dioxide, copovidone, ferric oxide, hypromellose, mannitol, medium-chain triglycerides, microcrystalline cellulose, natural and artificial flavor, pharmaceutical ink, polyethylene glycol, propylene glycol, sodium lauryl sulfate, sucralose, titanium dioxide

***Questions or comments?*** Call weekdays 9 AM to 5 PM EST at **1-800-88-ADVIL**