

Anbesol[®] Maximum Strength Liquid

This is the most current labeling information, and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts

Active ingredient

Benzocaine 20% Oral anesthetic

Purpose

Uses

- temporarily relieves pain associated with the following mouth and gum irritations:
 - toothache
 - sore gums
 - canker sores
 - braces
 - minor dental procedures
 - dentures

Warnings

METHEMOGLOBINEMIA WARNING: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics

Do not use

- for teething
- in children under age 2

When using this product

- avoid contact with the eyes
- do not exceed recommended dosage
- do not use for more than 7 days unless directed by a doctor/dentist

Stop use and ask a doctor if

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling, rash, or fever develops

Keep out of reach of children. If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

Anbesol[®] Maximum Strength Liquid

This is the most current labeling information, and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Directions

- adults and children 2 years of age and older:
 - wipe liquid on with cotton, or cotton swab, or fingertip
 - apply to the affected area up to 4 times daily or as directed by a doctor/dentist
- children under 12 years of age: adult supervision should be given in the use of this product
- children under 2 years of age: do not use

Other information store at 20-25°C (68-77°F)

Inactive ingredients benzyl alcohol, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, methylparaben, natural flavor, polyethylene glycol, polysorbate 80, propylene glycol, saccharin

Questions or comments? Call weekdays 9 AM to 5 PM EST at
1-888-797-5638