

# SUCCINYLCHOLINE CHLORIDE



QUELICIN®

20 mg/mL Solution for Injection (IM/IV)


## 1.0 PHARMACOLOGIC CATEGORY

Neuromuscular blocker

## 2.0 DESCRIPTION

Succinylcholine Chloride (Quelicin) is a sterile, nonpyrogenic solution to be used as a short-acting, depolarizing neuromuscular blocker for intravenous or intramuscular use. Succinylcholine Chloride (Quelicin) contains succinylcholine chloride as the active pharmaceutical ingredient.

Succinylcholine Chloride is chemically designated  $C_{14}H_{30}Cl_2N_2O_4$  and its molecular weight is 361.31. The chemical name of succinylcholine chloride is ethanaminium, 2,2'- [(1,4-dioxo-1,4 butanediyl) bis(oxy)] bis[N,N,N-trimethyl-], dichloride. It has the following structural formula:



## 3.0 FORMULATION

Each mL contains Succinylcholine Chloride, 20 mg.

Succinylcholine chloride is a diquaternary base consisting of the dichloride salt of the dicholine ester of succinic acid. It is a white, odorless, slightly bitter powder, very soluble in water.

Succinylcholine Chloride (Quelicin) 200 mg/10 mL (20 mg/mL) is intended for multiple dose administration and contains preservative. Each 1 mL of Succinylcholine Chloride (Quelicin) 200 mg/10 mL (20 mg/mL) multiple-dose fliptop vials contains: 20 mg of succinylcholine anhydrous (equivalent to 22.65 mg of Succinylcholine Chloride, USP), 1.8 mg of methylparaben and 0.2 mg of propylparaben as preservatives, 4.65 mg of sodium chloride as iso-osmotic agent, and sodium hydroxide and hydrochloric acid as pH adjusters in water for injection. The pH of the solution is between 3.0 and 4.5, with an osmolarity of 0.338 mOsm/mL (calc.).

## 4.0 CLINICAL PARTICULARS

### 4.1 Therapeutic Indications

Succinylcholine Chloride (Quelicin) is indicated in adults and pediatric patients:

- as an adjunct to general anesthesia
- to facilitate tracheal intubation
- to provide skeletal muscle relaxation during surgery or mechanical ventilation.

### 4.2 Dosage and Method of Administration

#### Important Dosage and Administration Information

- Succinylcholine Chloride (Quelicin) is for intravenous or intramuscular use only.
- Succinylcholine Chloride (Quelicin) must be titrated to effect by or under supervision of experienced clinicians who are familiar with its actions and with appropriate neuromuscular monitoring techniques.
- Succinylcholine Chloride (Quelicin) should be administered only by those skilled in the management of artificial respiration and only when facilities are instantly available for tracheal intubation and for providing adequate ventilation of the patient, including the administration of oxygen under positive pressure and the elimination of CO<sub>2</sub>. The clinician must be prepared to assist or control respiration.
- The dosage of Succinylcholine Chloride (Quelicin) should be individualized and should always be determined by the clinician after careful assessment of the patient.
- To avoid distress to the patient, do not administer Succinylcholine Chloride (Quelicin) before unconsciousness has been induced (see **section 4.4 Special Warnings and Precautions for Use**).
- The occurrence of bradyarrhythmias with administration of Succinylcholine Chloride (Quelicin) may be reduced by pretreatment with anticholinergics (e.g., atropine) (see **section 4.4 Special Warnings and Precautions for Use**).
- Monitor neuromuscular function with a peripheral nerve stimulator when using Succinylcholine Chloride (Quelicin) by infusion (see **sections 4.2 Dosage and Method of Administration** and **4.4 Special Warnings and Precautions for Use**).
- Visually inspect Succinylcholine Chloride (Quelicin) for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not administer solutions that are not clear and colorless.
- Succinylcholine Chloride (Quelicin) supplied in multiple dose vials does not require dilution before use (see **section 4.2 Dosage and Method of Administration**).

### **Risk of Medication Errors**

Accidental administration of neuromuscular blocking agents may be fatal. Store Succinylcholine Chloride (Quelicin) with the cap and ferrule intact and in a manner that minimizes the possibility of selecting the wrong product (see **section 4.4 Special Warnings and Precautions for Use**).

### **Dosage Recommendations for Intravenous Use in Adults**

#### **For Short Surgical Procedures**

The average dose required to produce neuromuscular blockade and to facilitate tracheal intubation is 0.6 mg/kg Succinylcholine Chloride (Quelicin) given intravenously. The optimum intravenous dose of Succinylcholine Chloride (Quelicin) will vary among patients and may be from 0.3 mg/kg to 1.1 mg/kg for adults. Following intravenous administration of doses in this range, neuromuscular blockade develops in about 1 minute; maximum blockade may persist for about 2 minutes, after which recovery takes place within 4 to 6 minutes. A 5 to 10 mg intravenous test dose of Succinylcholine Chloride (Quelicin) may be used to determine the sensitivity of the patient and the individual recovery time (see **section 4.4 Special Warnings and Precautions for Use**).

#### **For Long Surgical Procedures**

##### *Continuous Intravenous Infusion*

The dosage of Succinylcholine Chloride (Quelicin) administered by continuous intravenous infusion depends upon the duration of the surgical procedure and the need for muscle relaxation.

Diluted Succinylcholine Chloride (Quelicin) solutions containing from 1 mg/mL to 2 mg/mL succinylcholine have commonly been used for continuous intravenous infusion (see **section 4.2 Dosage and Method of Administration**). The more dilute solution (1 mg/mL) is probably preferable from the standpoint of ease of control of the rate of administration of Succinylcholine Chloride (Quelicin) and, hence, of relaxation. This diluted Succinylcholine Chloride (Quelicin) solution containing 1 mg/mL succinylcholine may be administered intravenously at a rate of 0.5 mg (0.5 mL) per minute to 10 mg (10 mL) per minute to obtain the required amount of relaxation. The amount required per minute will depend upon the individual response as well as the degree of relaxation required. The average rate of continuous intravenous infusion for an adult ranges between 2.5 mg per minute and 4.3 mg per minute.

Monitor neuromuscular function with a peripheral nerve stimulator when using Succinylcholine Chloride (Quelicin) by infusion in order to avoid overdose, detect development of Phase II block, follow its rate of recovery, and assess the effects of reversing agents (see **section 4.4 Special Warnings and Precautions for Use**).

##### *Intermittent Intravenous Injection*

Intermittent intravenous injections of Succinylcholine Chloride (Quelicin) may also be used to provide muscle relaxation for long procedures. An intravenous injection of 0.3 mg/kg to 1.1 mg/kg may be given initially, followed, at appropriate intervals, by further intravenous injections of 0.04 mg/kg to 0.07 mg/kg to maintain the degree of relaxation required.

## **Dosage Recommendations for Intravenous Use in Pediatric Patients**

For emergency tracheal intubation or in instances where immediate securing of the airway is necessary, the intravenous dose of Succinylcholine Chloride (Quelicin) is 2 mg/kg for infants and other small pediatric patients; for older pediatric patients and adolescents the intravenous dose is 1 mg/kg (see **section 4.4 Special Warnings and Precautions for Use** and **4.6 Fertility, Pregnancy and Lactation**). The effective dose of Succinylcholine Chloride (Quelicin) in pediatric patients may be higher than that predicted by body weight dosing alone. For example, the usual adult intravenous dose of 0.6 mg/kg is comparable to a dose of 2 mg/kg to 3 mg/kg in neonates and infants up to 6 months of age and 1 mg/kg to 2 mg/kg in infants up to 2 years of age (see **section 5.2 Pharmacokinetic Properties**).

## **Dosage Recommendations for Intramuscular Use in Adults and Pediatric Patients**

If a suitable vein is inaccessible, Succinylcholine Chloride (Quelicin) may be administered intramuscularly at a dose of up to 3 mg/kg to 4 mg/kg to infants, older pediatric patients or adults. The total dose administered by the intramuscular route should not exceed 150 mg. The onset of effect of succinylcholine given intramuscularly is usually observed in about 2 to 3 minutes.

### **Drug compatibility**

Succinylcholine Chloride (Quelicin) is acidic (pH is between 3.0 and 4.5) and may not be compatible with alkaline solutions having a pH greater than 8.5 (e.g., barbiturate solutions). Therefore, do not mix Succinylcholine Chloride (Quelicin) with alkaline solutions.

## **Preparation of Succinylcholine Chloride (Quelicin)**

Succinylcholine Chloride (Quelicin) supplied in multiple-dose vials does not require dilution before use.

Succinylcholine Chloride (Quelicin) may be diluted to 1 mg/mL or 2 mg/mL in a solution such as:

- 5% Dextrose Injection, USP, or
- 0.9% Sodium Chloride Injection, USP

Prepare the diluted Succinylcholine Chloride (Quelicin) solution for single patient use only. Store the diluted Succinylcholine Chloride (Quelicin) solution in a refrigerator [2 °C to 8 °C (36 °F to 46 °F)] and use within 24 hours after preparation. Visually inspect the diluted Succinylcholine Chloride (Quelicin) solution for particulate matter and discoloration prior to administration. Do not administer solutions that are not clear and colorless. Discard any unused portion of the diluted Succinylcholine Chloride (Quelicin) solution.

## **4.3 Contraindications**

Succinylcholine Chloride (Quelicin) is contraindicated:

- in patients with skeletal muscle myopathies (see **section 4.4 Special Warnings and Precautions for Use**)

- in patients with known hypersensitivity to succinylcholine. Severe anaphylactic reactions to succinylcholine have been reported (see **section 4.4 Special Warnings and Precautions for Use**)
- after the acute phase of injury following major burns, multiple trauma, extensive denervation of skeletal muscle, or upper motor neuron injury, which may result in severe hyperkalemia and cardiac arrest (see **section 4.4 Special Warnings and Precautions for Use**).
- in patients with known or suspected genetic susceptibility to malignant hyperthermia (see **section 4.4 Special Warnings and Precautions for Use** and **section 5.1 Pharmacodynamic Properties - Pharmacogenomics**)

#### 4.4 Special Warnings and Precautions for Use

##### Warnings

##### **Ventricular dysrhythmias, cardiac arrest, and death from hyperkalemic rhabdomyolysis in pediatric patients**

- Acute rhabdomyolysis with hyperkalemia followed by ventricular dysrhythmias, cardiac arrest, and death has occurred after the administration of succinylcholine to apparently healthy pediatric patients who were subsequently found to have undiagnosed skeletal muscle myopathy, most frequently Duchenne muscular dystrophy (see **section 4.4 Special Warnings and Precautions for Use**).
- When a healthy appearing pediatric patient develops cardiac arrest within minutes after administration of Succinylcholine Chloride (Quelicin) , not felt to be due to inadequate ventilation, oxygenation or anesthetic overdose, immediate treatment for hyperkalemia should be instituted. In the presence of signs of malignant hyperthermia, appropriate treatment should be instituted concurrently (see **section 4.4 Special Warnings and Precautions for Use**).
- Reserve the use of Succinylcholine Chloride (Quelicin) in pediatric patients for emergency intubation or instances where immediate securing of the airway is necessary, e.g., laryngospasm, difficult airway, full stomach, or for intramuscular use when a suitable vein is inaccessible (see **section 4.4 Special Warnings and Precautions for Use**).

##### **Ventricular Dysrhythmias, Cardiac Arrest, and Death From Hyperkalemic Rhabdomyolysis in Pediatric Patients**

There have been reports of ventricular dysrhythmias, cardiac arrest, and death secondary to acute rhabdomyolysis with hyperkalemia in apparently healthy pediatric patients who received succinylcholine. Many of these pediatric patients were subsequently found to have a skeletal muscle myopathy such as Duchenne muscular dystrophy whose clinical signs were not obvious.

The syndrome often presented as sudden cardiac arrest within minutes after the administration of succinylcholine. These pediatric patients were usually, but not exclusively, males, and most frequently 8 years of age or younger. There have also

been reports in adolescents. There may be no signs or symptoms to alert the practitioner to which patients are at risk. A careful history and physical may identify developmental delays suggestive of a myopathy. A preoperative creatine kinase could identify some but not all patients at risk.

When a healthy appearing pediatric patient develops cardiac arrest within minutes after administration of Succinylcholine Chloride (Quelicin), not felt to be due to inadequate ventilation, oxygenation or anesthetic overdose, immediate treatment for hyperkalemia should be instituted. Due to the abrupt onset of this syndrome, routine resuscitative measures are likely to be unsuccessful. Careful monitoring of the electrocardiogram may alert the practitioner to peaked T waves (an early sign). Administration of intravenous calcium, bicarbonate, and glucose with insulin, with hyperventilation have resulted in successful resuscitation in some of the reported cases. Extraordinary and prolonged resuscitative efforts have been effective in some cases. In addition, in the presence of signs of malignant hyperthermia, appropriate treatment should be initiated concurrently (see **section 4.4 Special Warnings and Precautions for Use**).

Because it is difficult to identify which patients are at risk, reserve the use of Succinylcholine Chloride (Quelicin) in pediatric patients for emergency intubation or instances where immediate securing of the airway is necessary, e.g., laryngospasm, difficult airway, full stomach, or for intramuscular use when a suitable vein is inaccessible.

#### **Anaphylaxis**

Severe anaphylactic reactions to neuromuscular blocking agents, including succinylcholine, have been reported. These reactions have, in some cases, been life-threatening and fatal. Due to the potential severity of these reactions, the necessary precautions, such as the immediate availability of appropriate emergency treatment, should be taken. Allergic cross-reactivity between neuromuscular blocking agents, both depolarizing and non-depolarizing, has been reported in this class of drugs. Therefore, assess patients for previous anaphylactic reactions to other neuromuscular blocking agents before administering Succinylcholine Chloride (Quelicin).

#### **Risk of Death due to Medication Errors**

Administration of Succinylcholine Chloride (Quelicin) results in paralysis, which may lead to respiratory arrest and death; this progression may be more likely to occur in a patient for whom it is not intended. Confirm proper selection of intended product and avoid confusion with other injectable solutions that are present in critical care and other clinical settings. If another healthcare provider is administering the product, ensure that the intended dose is clearly labeled and communicated.

#### **Hyperkalemia**

Succinylcholine Chloride (Quelicin) may induce serious cardiac arrhythmias or cardiac arrest due to hyperkalemia in patients with electrolyte abnormalities and those who may have digitalis toxicity.

Succinylcholine Chloride (Quelicin) is contraindicated after the acute phase of injury following major burns, multiple trauma, extensive denervation of skeletal muscle, or upper motor neuron injury (see **section 4.3 Contraindications**). The risk of hyperkalemia in these patients increases over time and usually peaks at 7 to 10 days

after the injury. The risk is dependent on the extent and location of the injury. The precise time of onset and the duration of the risk period are undetermined.

Patients with chronic abdominal infection, subarachnoid hemorrhage, or conditions causing degeneration of central and peripheral nervous systems are at an increased risk of developing severe hyperkalemia after Succinylcholine Chloride (Quelicin) administration. Consider avoiding use of Succinylcholine Chloride (Quelicin) in these patients or verify the patient's baseline potassium levels are within the normal range prior to Succinylcholine Chloride (Quelicin) administration.

### **Malignant Hyperthermia**

In susceptible individuals, succinylcholine may trigger malignant hyperthermia, a skeletal muscle hypermetabolic state leading to high oxygen demand. Fatal outcomes of malignant hyperthermia have been reported.

The risk of developing malignant hyperthermia increases with the concomitant administration of succinylcholine and volatile anesthetic agents. Succinylcholine Chloride (Quelicin) can induce malignant hyperthermia in patients with known or suspected susceptibility based on genetic factors or family history, including those with certain inherited ryanodine receptor (RYR1) or dihydropyridine receptor (CACNA1S) variants (see **section 4.3 Contraindications and Pharmacogenomics**).

Signs consistent with malignant hyperthermia may include hyperthermia, hypoxia, hypercapnia, muscle rigidity (e.g., jaw muscle spasm), tachycardia (e.g., particularly that unresponsive to deepening anesthesia or analgesic medication administration), tachypnea, cyanosis, arrhythmias, hypovolemia, and hemodynamic instability. Skin mottling, coagulopathies, and renal failure may occur later in the course of the hypermetabolic process.

Successful treatment of malignant hyperthermia depends on early recognition of the clinical signs. If malignant hyperthermia is suspected, discontinue all triggering agents (i.e., volatile anesthetic agents and succinylcholine), administer intravenous dantrolene sodium, and initiate supportive therapies. Consult prescribing information for intravenous dantrolene sodium for additional information on patient management. Supportive therapies include administration of supplemental oxygen and respiratory support based on clinical need, maintenance of hemodynamic stability and adequate urinary output, management of fluid and electrolyte balance, correction of acid base derangements, and institution of measures to control rising temperature.

### **Bradycardia**

Intravenous bolus administration of Succinylcholine Chloride (Quelicin) in pediatric patients (including infants) may result in profound bradycardia or, rarely, asystole. In both adults and pediatric patients the incidence of bradycardia, which may progress to asystole, is higher following a second dose of succinylcholine. The incidence and severity of bradycardia is higher in pediatric patients than adults. Whereas bradycardia is common in pediatric patients after an initial dose of 1.5 mg/kg, bradycardia is seen in adults only after repeated exposure. Pretreatment with anticholinergic agents (e.g., atropine) may reduce the occurrence of bradyarrhythmias.

### **Increase in Intraocular Pressure**

Succinylcholine causes an increase in intraocular pressure. Avoid Succinylcholine Chloride (Quelicin) in instances in which an increase in intraocular pressure is undesirable (e.g., narrow angle glaucoma, penetrating eye injury) unless the potential benefit of its use outweighs the potential risk.

### **Prolonged Neuromuscular Block due to Phase II Block and Tachyphylaxis**

When Succinylcholine Chloride (Quelicin) is given over a prolonged period of time, the characteristic depolarization block of the myoneural junction (Phase I block) may change to a block with characteristics superficially resembling a non-depolarizing block (Phase II block). Prolonged respiratory muscle paralysis or weakness may be observed in patients manifesting this transition to Phase II block. Tachyphylaxis occurs with repeated administration (see **section 5.1 Pharmacodynamic Properties**).

When Phase II block is suspected in cases of prolonged neuromuscular blockade, positive diagnosis should be made by peripheral nerve stimulation, prior to administration of any anticholinesterase drug. Reversal of Phase II block is a medical decision which must be made upon the basis of the patient, clinical pharmacology, and the experience and judgment of the clinician. The presence of Phase II block is indicated by fade of responses to successive stimuli (preferably "train of four"). The use of an anticholinesterase drug such as neostigmine to reverse Phase II block should be accompanied by appropriate doses of an anticholinergic drug to prevent disturbances of cardiac rhythm. After adequate reversal of Phase II block with an anticholinesterase agent, the patient should be continually observed for at least 1 hour for signs of return of muscle relaxation. Reversal should not be attempted unless: (1) a peripheral nerve stimulator is used to determine the presence of Phase II block (since anticholinesterase agents will potentiate succinylcholine-induced Phase I block), and (2) spontaneous recovery of muscle twitch has been observed for at least 20 minutes and has reached a plateau with further recovery proceeding slowly; this delay is to ensure complete hydrolysis of succinylcholine by plasma cholinesterase prior to administration of the anticholinesterase agent. Should the type of block be misdiagnosed, depolarization of the type initially induced by succinylcholine (i.e., Phase I block) will be prolonged by an anticholinesterase agent.

### **Risk of Prolonged Neuromuscular Block in Patients with Reduced Plasma Cholinesterase Activity**

Succinylcholine Chloride (Quelicin) is not recommended in patients with known reduced plasma cholinesterase (pseudocholinesterase) activity due to the likelihood of prolonged neuromuscular block following administration of Succinylcholine Chloride (Quelicin) in such patients.

Plasma cholinesterase activity may be diminished in the presence of genetic abnormalities of plasma cholinesterase (e.g., patients heterozygous or homozygous for atypical plasma cholinesterase gene), pregnancy, severe liver or kidney disease, malignant tumors, infections, burns, anemia, decompensated heart disease, peptic ulcer, or myxedema. Plasma cholinesterase activity may also be diminished by chronic administration of oral contraceptives, glucocorticoids, or certain monoamine oxidase inhibitors and by irreversible inhibitors of plasma cholinesterase (e.g., organophosphate insecticides, echothiophate, and certain antineoplastic drugs) [(see **section 4.5 Interaction with other medicinal products and other forms of Interaction**)].

Patients homozygous for atypical plasma cholinesterase gene (1 in 2,500 patients) are extremely sensitive to the neuromuscular blocking effect of succinylcholine. If Succinylcholine Chloride (Quelicin) is administered to a patient homozygous for atypical plasma cholinesterase, resulting apnea or prolonged muscle paralysis should be treated with controlled respiration.

**Risk of Additional Trauma in Patients with Fractures or Muscle Spasms**

Succinylcholine should be employed with caution in patients with fractures or muscle spasm because the initial muscle fasciculations may cause additional trauma.

**Increase in Intracranial Pressure**

Succinylcholine Chloride (Quelicin) may cause a transient increase in intracranial pressure; however, adequate anesthetic induction prior to administration of Succinylcholine Chloride (Quelicin) will minimize this effect.

**Risk of Aspiration due to Increase in Intra-gastric Pressure**

Succinylcholine may increase intra-gastric pressure, which could result in regurgitation and possible aspiration of stomach contents. Evaluate patients at risk for aspiration and regurgitation. Monitor patients during induction of anesthesia and neuromuscular blockade for clinical signs of vomiting and/or aspiration.

**Prolonged Neuromuscular Block in Patients with Hypokalemia or Hypocalcemia**

Neuromuscular blockade may be prolonged in patients with hypokalemia (e.g., after severe vomiting, diarrhea, digitalization and diuretic therapy) or hypocalcemia (e.g., after massive transfusions). Correct severe electrolyte disturbances when possible. In order to help preclude possible prolongation of neuromuscular block, monitor neuromuscular transmission throughout the use of Succinylcholine Chloride (Quelicin).

### **Risks due to Inadequate Anesthesia**

Neuromuscular blockade in the conscious patient can lead to distress. Use Succinylcholine Chloride (Quelicin) in the presence of appropriate sedation or general anesthesia. Monitor patients to ensure that the level of anesthesia is adequate. In emergency situations, however, it may be necessary to administer Succinylcholine Chloride (Quelicin) before unconsciousness is induced.

### **Pediatric Use**

Safety and effectiveness of Succinylcholine Chloride have been established in pediatric patient age groups, neonate to adolescent. Because of a risk of ventricular dysrhythmias, cardiac arrest, and death from hyperkalemic rhabdomyolysis in pediatric patients, reserve the use of Succinylcholine Chloride (Quelicin) in pediatric patients for emergency intubation or instances where immediate securing of the airway is necessary, e.g., laryngospasm, difficult airway, full stomach, or for intramuscular use when a suitable vein is inaccessible (see **section 4.4 Special Warnings and Precautions for Use**).

Intravenous bolus administration of Succinylcholine Chloride (Quelicin) in pediatric patients (including infants) may result in profound bradycardia or, rarely, asystole. The incidence and severity of bradycardia is higher in pediatric patients than adults.

The effective dose of Succinylcholine Chloride (Quelicin) in pediatric patients may be higher than that predicted by body weight dosing alone (see **section 4.2 Dosage and Administration**).

### **Geriatric Use**

Clinical studies of Succinylcholine Chloride (Quelicin) did not include sufficient numbers of subjects aged 65 years and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

## **4.5 Interaction with other medicinal products and other forms of Interaction**

Drugs that may enhance the neuromuscular blocking action of succinylcholine include: promazine, oxytocin, aprotinin, certain non-penicillin antibiotics, quinidine,  $\beta$ -adrenergic blockers, procainamide, lidocaine, trimethaphan, lithium carbonate, magnesium salts, quinine, chloroquine, isoflurane, desflurane, metoclopramide and terbutaline.

The neuromuscular blocking effect of succinylcholine may be enhanced by drugs that reduce plasma cholinesterase activity (e.g., chronically administered oral contraceptives, glucocorticoids, or certain monoamine oxidase inhibitors) or by drugs that irreversibly inhibit plasma cholinesterase (see **section 4.4 Special Warnings and Precautions for Use**).

If other neuromuscular blocking agents are to be used during the same procedure, consider the possibility of a synergistic or antagonistic effect.

## 4.6 Fertility, Pregnancy and Lactation

### *Pregnancy*

#### **Risk Summary**

Available data from published literature from case reports and case series over decades of use with succinylcholine during pregnancy have not identified a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. Succinylcholine is used commonly during delivery by caesarean section to provide muscle relaxation. If succinylcholine is used during labor and delivery, there is a risk for prolonged apnea in some pregnant women (see **Clinical Considerations**). Animal reproduction studies have not been conducted with Succinylcholine Chloride.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

#### **Clinical Considerations**

##### *Maternal Adverse Reactions*

Plasma cholinesterase levels are decreased by approximately 24% during pregnancy and for several days postpartum which can prolong the effect of succinylcholine. Therefore, some pregnant patients may experience prolonged apnea.

##### *Fetal/Neonatal Adverse Reactions*

Apnea and flaccidity may occur in the newborn after repeated high doses to, or in the presence of atypical plasma cholinesterase in, the mother.

##### *Labor or Delivery*

Succinylcholine is commonly used to provide muscle relaxation during delivery by caesarean section. Succinylcholine is known to cross the placental barrier in an amount that is dependent on the concentration gradient between the maternal and fetal circulation.

### *Lactation*

#### **Risk Summary**

There are no data on the presence of succinylcholine or its metabolite in either human or animal milk, the effects on the breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Succinylcholine Chloride (Quelicin) and any potential adverse effects on the breastfed infant from Succinylcholine Chloride (Quelicin) or from the underlying maternal condition.

## 4.7 Effects on Ability to Drive and Use Medicines

No information available.

## 4.8 Undesirable Effects

The following clinically significant adverse reactions are discussed in greater detail in other sections of the labeling:

- Ventricular Dysrhythmias, Cardiac Arrest, and Death from Hyperkalemic Rhabdomyolysis in Pediatric Patients (see **section 4.4 Special Warnings and Precautions for Use**)
- Anaphylaxis (see **section 4.4 Special Warnings and Precautions for Use**)
- Hyperkalemia (see **section 4.4 Special Warnings and Precautions for Use**)
- Malignant Hyperthermia (see **section 4.4 Special Warnings and Precautions for Use**)
- Bradycardia (see **section 4.4 Special Warnings and Precautions for Use**)
- Increase in Intraocular Pressure (see **section 4.4 Special Warnings and Precautions for Use**)
- Prolonged Neuromuscular Block due to Phase II Block and Tachyphylaxis (see **section 4.4 Special Warnings and Precautions for Use**)

The following adverse reactions associated with the use of succinylcholine were identified in clinical studies or postmarketing reports. Because some of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure:

Cardiovascular disorders: Cardiac arrest, arrhythmias, bradycardia, tachycardia, hypertension, hypotension

Electrolyte disorders: Hyperkalemia

Eye disorders: Increased intraocular pressure

Gastrointestinal disorders: Excessive salivation

Immune system disorders: Hypersensitivity reactions including anaphylaxis (in some cases life-threatening and fatal)

Musculoskeletal disorders: Malignant hyperthermia, rhabdomyolysis with possible myoglobinuric acute renal failure, muscle fasciculation, jaw rigidity, postoperative muscle pain

Respiratory disorders: Prolonged respiratory depression or apnea

Skin disorders: Rash

#### **4.9 Overdose and Treatment**

Overdosage with Succinylcholine Chloride (Quelicin) may result in neuromuscular block beyond the time needed for surgery and anesthesia. This may be manifested by skeletal muscle weakness, decreased respiratory reserve, low tidal volume, or apnea. The primary treatment is maintenance of a patent airway and respiratory support until recovery of normal respiration is assured. Depending on the dose and duration of Succinylcholine Chloride (Quelicin) administration, the characteristic depolarizing neuromuscular block (Phase I) may change to a block with characteristics superficially resembling a non-depolarizing block (Phase II) (see **section 4.4 Special Warnings and Precautions for Use**).

### **5.0 PHARMACOLOGICAL PROPERTIES**

#### **Mechanism of Action**

Succinylcholine is a depolarizing neuromuscular blocker. As does acetylcholine, it combines with the cholinergic receptors of the motor end plate to produce depolarization. This depolarization may be observed as fasciculations. Subsequent neuromuscular transmission is inhibited so long as adequate concentration of succinylcholine remains at the receptor site. Onset of flaccid paralysis is rapid (less than one minute after intravenous administration) and with single administration lasts approximately 4 to 6 minutes.

The paralysis following administration of succinylcholine is progressive, with differing sensitivities of different muscles. This initially involves consecutively the levator muscles of the face, muscles of the glottis and finally the intercostals and the diaphragm and all other skeletal muscles.

### **5.1 Pharmacodynamic Properties**

Depending on the dose and duration of succinylcholine administration, the characteristic depolarizing neuromuscular block (Phase I block) may change to a block with characteristics superficially resembling a non depolarizing block (Phase II block). This may be associated with prolonged respiratory muscle paralysis or weakness in patients who manifest the transition to Phase II block. Tachyphylaxis occurs with repeated administration (see **section 4.4 Special Warnings and Precautions for Use**). The transition from Phase I to Phase II block has been reported in 7 of 7 patients studied under halothane anesthesia after an accumulated dose of 2 to 4 mg/kg succinylcholine (administered in repeated, divided doses). The onset of Phase II block coincided with the onset of tachyphylaxis and prolongation of spontaneous recovery. In another study, using balanced anesthesia (N<sub>2</sub>O/O<sub>2</sub>/narcotic thiopental) and succinylcholine infusion, the transition was less abrupt, with great individual variability in the dose of succinylcholine required to produce Phase II block. Of 32 patients studied, 24 developed Phase II block. Tachyphylaxis was not associated with the transition to Phase II block, and 50% of the patients who developed Phase II block experienced prolonged recovery (see **section 4.4 Special Warnings and Precautions for Use**).

Succinylcholine has no direct effect on the myocardium. Succinylcholine stimulates both autonomic ganglia and muscarinic receptors which may cause changes in cardiac rhythm, including cardiac arrest. Changes in rhythm, including cardiac arrest, may also result from vagal stimulation, which may occur during surgical procedures, or from hyperkalemia, particularly in pediatric patients (see **sections 4.4 Special Warnings and Precautions for Use** and **4.6 Fertility, Pregnancy and Lactation**). These effects are enhanced by halogenated anesthetics.

Succinylcholine causes an increase in intraocular pressure immediately after its injection and during the fasciculation phase, and increases which may persist after onset of complete paralysis (see **section 4.4 Special Warnings and Precautions for Use**).

Succinylcholine may cause increases in intracranial pressure immediately after its injection and during the fasciculation phase (see **section 4.4 Special Warnings and Precautions for Use**).

As with other neuromuscular blocking agents, the potential for releasing histamine is present following succinylcholine administration. Signs and symptoms of histamine mediated release such as flushing, hypotension and bronchoconstriction are, however, uncommon with normal clinical usage.

Succinylcholine has no effect on consciousness, pain threshold or cerebation (see **section 4.4 Special Warnings and Precautions for Use**).

Succinylcholine has no direct action on the uterus or other smooth muscle structures.

### **Pharmacogenomics**

RYR1 and CACNA1S are polymorphic genes and multiple pathogenic variants have been associated with malignant hyperthermia susceptibility (MHS) in patients receiving succinylcholine, including Succinylcholine Chloride (Quelicin). Case reports as well as exvivo studies have identified multiple variants in RYR1 and CACNA1S associated with MHS. Variant pathogenicity should be assessed based on prior clinical experience, functional studies, prevalence information, or other evidence (see **sections 4.3 Contraindications and 4.4 Special Warnings and Precautions for Use**).

## **5.2 Pharmacokinetic Properties**

### Elimination

Succinylcholine levels were reported to be below the detection limit of 2 µg/mL after 2.5 minutes of an intravenous bolus dose of 1 or 2 mg/kg in 14 anesthetized patients.

### Metabolism

Succinylcholine is rapidly hydrolyzed by plasma cholinesterase to succinylmonocholine (which possesses clinically insignificant depolarizing muscle relaxant properties) and then more slowly to succinic acid and choline.

### Excretion

About 10% of the drug is excreted unchanged in the urine.

### Specific Populations

#### *Pediatric Patients*

Due to the relatively large volume of distribution in the pediatric patient versus the adult patient, the effective dose of Succinylcholine Chloride (Quelicin) in pediatric patients may be higher than that predicted by body weight dosing alone (see **section 4.2 Dosage and Method of Administration**).

## **5.3 Preclinical Safety Data**

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### Carcinogenesis

There have been no long-term studies performed in animals to evaluate carcinogenic potential of succinylcholine.

### Mutagenesis

Adequate studies have not been completed to evaluate the genotoxic potential of succinylcholine.

### Impairment of Fertility

There are no studies to evaluate the potential impact of succinylcholine on fertility.

## **6.0 PHARMACEUTICAL PARTICULARS**

### **6.1 Shelf-Life**

15 Months

### **6.2 Storage Condition**

Store at 2°C to 8°C (36°F to 46°F). The multiple-dose vials are stable for up to 14 days at room temperature without significant loss of potency.

### **6.3 Availability**

Quelicin Solution for Injection (IM/IV) is available in 20 mg/mL in clear glass vial USP Type I with gray rubber butyl stopper and sealed with red aluminum and red flip-off plastic cap 1 x 10 mL.

## **7.0 FDA REGISTRATION NUMBER**

Quelicin 20 mg/mL Solution for Injection (IM/IV): DR-699

## **8.0 DATE OF FIRST AUTHORIZATION**

Quelicin 20 mg/mL Solution for Injection (IM/IV): 22-Apr-1992

Keep out of reach of children.

**For suspected adverse drug reaction, report to the FDA: [www.fda.gov.ph](http://www.fda.gov.ph)**

**Seek medical attention immediately at the first sign of any adverse drug reaction.**

**CAUTION:** Foods, Drugs, Devices, and Cosmetics Act prohibits dispensing without prescription.

### **Manufactured by:**

Hospira Inc, Highway 301,  
North, Rocky Mount,  
North Carolina, USA

### **Marketing Authorization Holder:**

Pfizer, Inc.  
19F-20F, 8 Rockwell Building, Hidalgo Drive,  
Rockwell Center, Poblacion,

Makati City 1210 Metro Manila, Philippines

Revision No: 1.1 Revision Date: 04 March 2024 Reference: USPI, Ver.No: LAB-1246-5.0 + BOH request Reference Date: 19 September 2022
---