

Voriconazole Tablets I.P. and Voriconazole Powder for Solution for Infusion I.P. VFEND™



1. GENERIC NAME

Voriconazole Tablets I.P. and Voriconazole Powder for Solution for Infusion I.P.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Tablets:

Each film-coated tablet contains 50 mg or 200 mg voriconazole.

Powder for solution for infusion:

Each vial contains voriconazole 200 mg equivalent to 10 mg/ml solution following reconstitution (see section 8.4).

List of Excipients

Film-coated tablets:

Tablet Core:

Lactose Monohydrate
Pregelatinised Starch
Croscarmellose Sodium
Povidone
Magnesium Stearate

Film-coating:

Hypromellose
Titanium Dioxide (E171)
Lactose Monohydrate
Glycerol Triacetate (Triacetin)

Powder for solution for infusion:

Sulfobutylether Beta Cyclodextrin Sodium
Water for Injections

All strengths/presentations mentioned in this document might not be available in the market.

™ Trademark Proprietor: Pfizer Inc., USA
Licensed User: Pfizer Products India Private Limited, India

3. DOSAGE FORM AND STRENGTH

Tablets:

Voriconazole 50 mg film-coated tablets are white, round tablets, debossed “Pfizer” on one side and “VOR50” on the reverse (tablets).

Voriconazole 200 mg film-coated tablets are white, capsule-shaped tablets, debossed “Pfizer” on one side and “VOR200” on the reverse (tablets).

Powder for solution for infusion:

Voriconazole powder for solution for infusion is a white lyophilised powder containing nominally 200 mg voriconazole presented in a 30 ml clear glass vial.

4. CLINICAL PARTICULARS

4.1 Therapeutic indication

Voriconazole is a broad spectrum, triazole antifungal agent and is indicated as follows:

Treatment of invasive aspergillosis;

Treatment of fluconazole-resistant serious invasive *Candida* infections (including *C. krusei*);

Treatment of oesophageal candidiasis;

Treatment of serious fungal infection caused by *Scedosporium* spp. and *Fusarium* spp. including *Fusarium solani* in patients intolerant of or refractory to other therapy.

Candidemia in non-neutropenic patients and the following *Candida* infections: disseminated infections in skin and infections in abdomen, kidney, bladder wall, and wounds.

4.2 Posology and method of administration

Posology

Electrolyte disturbances such as hypokalaemia, hypomagnesaemia and hypocalcaemia should be monitored and corrected, if necessary, prior to initiation and during voriconazole therapy (see section 4.4).

Powder for solution for infusion:

It is recommended that voriconazole is administered at a maximum rate of 3 mg/kg per hour over 1 to 3 hours.

Treatment

Adults

Therapy must be initiated with the specified loading dose regimen of either intravenous or oral voriconazole to achieve plasma concentrations on Day 1 that are close to steady state. On the basis

of the high oral bioavailability (96%; see section 5.3), switching between intravenous and oral administration is appropriate when clinically indicated.

Detailed information on dosage recommendations is provided in the following table:

	Intravenous	Oral	
		Patients 40 kg and above*	Patients less than 40 kg*
Loading dose regimen (first 24 hours)	6 mg/kg every 12 hours	400 mg every 12 hours	200 mg every 12 hours
Maintenance dose (after first 24 hours)	4 mg/kg twice daily	200 mg twice daily	100 mg twice daily

* This also applies to patients aged 15 years and older.

Duration of treatment

Treatment duration should be as short as possible depending on the patient's clinical and mycological response. Long term exposure to voriconazole greater than 180 days (6 months) requires careful assessment of the benefit-risk balance (see sections 4.4 and 5.2).

Dosage adjustment (Adults)

If patient response to treatment is inadequate, the maintenance dose may be increased to 300 mg twice daily for oral administration. For patients less than 40 kg the oral dose may be increased to 150 mg twice daily.

Tablets:

If patient is unable to tolerate treatment at a higher dose, reduce the oral dose by 50 mg steps to the 200 mg twice daily (or 100 mg twice daily for patients less than 40 kg) maintenance dose.

Powder for solution for infusion:

If patient is unable to tolerate intravenous treatment at 4 mg/kg twice daily, reduce the dose to 3 mg/kg twice daily.

Children (2 to <12 years) and young adolescents with low body weight (12 to 14 years and <50 kg)

Voriconazole should be dosed as children as these young adolescents may metabolise voriconazole more similarly to children than to adults.

The recommended dosing regimen is as follows:

	Intravenous	Oral
Loading Dose Regimen (first 24 hours)	9 mg/kg every 12 hours	Not recommended
Maintenance Dose (after first 24 hours)	8 mg/kg twice daily	9 mg/kg twice daily (a maximum dose of 350 mg twice daily)

Note: Based on a population pharmacokinetic analysis in 112 immunocompromised paediatric patients aged 2 to <12 years and 26 immunocompromised adolescents aged 12 to <17 years.

It is recommended to initiate the therapy with intravenous regimen, and oral regimen should be considered only after there is a significant clinical improvement. It should be noted that an 8 mg/kg intravenous dose will provide voriconazole exposure approximately 2-fold higher than a 9 mg/kg oral dose.

Tablets:

These oral dose recommendations for children are based on studies in which voriconazole was administered as the powder for oral suspension. Bioequivalence between the powder for oral suspension and tablets has not been investigated in a paediatric population. Considering the assumed limited gastro-enteric transit time in paediatric patients, the absorption of tablets may be different in paediatric compared to adult patients. It is therefore recommended to use the oral suspension formulation in children aged 2 to <12.

All other adolescents (12 to 14 years and ≥ 50 kg; 15 to 17 years regardless of body weight)
Voriconazole should be dosed as adults.

Dosage adjustment (Children [2 to <12 years] and young adolescents with low body weight [12 to 14 years and <50 kg])

Tablets:

If patient response to treatment is inadequate, the dose may be increased by 1 mg/kg steps (or by 50 mg steps if the maximum oral dose of 350 mg was used initially). If patient is unable to tolerate treatment, reduce the dose by 1 mg/kg steps (or by 50 mg steps if the maximum oral dose of 350 mg was used initially).

Powder for solution for infusion:

If patient response to treatment is inadequate, the intravenous dose may be increased by 1 mg/kg steps. If patient is unable to tolerate treatment, reduce the intravenous dose by 1 mg/kg steps.

Use in paediatric patients aged 2 to <12 years with hepatic or renal insufficiency has not been studied (see sections 4.8 and 5.3).

The following instructions apply to treatment:

Dosage adjustment

In the case of treatment-related adverse events, discontinuation of voriconazole and use of alternative antifungal agents must be considered (see sections 4.4 and 4.8)

Dosage adjustments in case of coadministration

Phenytoin may be coadministered with voriconazole if the maintenance dose of voriconazole is increased from 200 mg to 400 mg orally, twice daily (100 mg to 200 mg orally, twice daily in patients less than 40 kg), see sections 4.4 and 4.5.

Powder for solution for infusion

Rifabutin or phenytoin may be coadministered with voriconazole if the maintenance dose of voriconazole is increased to 5 mg/kg intravenously twice daily, see sections 4.4 and 4.5.

The combination of voriconazole with rifabutin should, if possible be avoided. However, if the combination is strictly needed, the maintenance dose of voriconazole may be increased from 200 mg to 350 mg orally, twice daily (100 mg to 200 mg orally, twice daily in patients less than 40 kg), see sections 4.4 and 4.5.

Efavirenz may be coadministered with voriconazole if the maintenance dose of voriconazole is increased to 400 mg every 12 hours and the efavirenz dose is reduced by 50%, i.e., to 300 mg once daily. When treatment with voriconazole is stopped, the initial dosage of efavirenz should be restored (see sections 4.4 and 4.5).

Elderly

No dose adjustment is necessary for elderly patients (see section 5.3).

Renal impairment

Tablets:

The pharmacokinetics of orally administered voriconazole are not affected by renal impairment. Therefore, no adjustment is necessary for oral dosing for patients with mild to severe renal impairment (see section 5.3).

Powder for solution for infusion:

In patients with moderate to severe renal dysfunction (creatinine clearance <50 ml/min), accumulation of the intravenous vehicle, SBECD, occurs. Oral voriconazole should be administered to these patients, unless an assessment of the risk benefit to the patient justifies the use of intravenous voriconazole. Serum creatinine levels should be closely monitored in these patients and, if it increases, consideration should be given to changing to oral voriconazole therapy (see section 5.3).

Voriconazole is haemodialysed with a clearance of 121 ml/min. A 4-hour haemodialysis session does not remove a sufficient amount of voriconazole to warrant dose adjustment.

The intravenous vehicle, SBECD, is haemodialysed with a clearance of 55 ml/min.

Hepatic impairment

It is recommended that the standard loading dose regimens be used but that the maintenance dose be halved in patients with mild to moderate hepatic cirrhosis (Child-Pugh A and B) receiving voriconazole (see section 5.3).

Voriconazole has not been studied in patients with severe chronic hepatic cirrhosis (Child-Pugh C).

There is limited data on the safety of voriconazole in patients with abnormal liver function tests (aspartate transaminase [AST], alanine transaminase [ALT], alkaline phosphatase [ALP], or total bilirubin >5 times the upper limit of normal).

Voriconazole has been associated with elevations in liver function tests and clinical signs of liver damage, such as jaundice, and must only be used in patients with severe hepatic impairment if the benefit outweighs the potential risk. Patients with severe hepatic impairment must be carefully monitored for drug toxicity (see section 4.8).

Paediatric population

The safety and efficacy of voriconazole in children below 2 years has not been established. Currently available data are described in sections 4.8 and 5.2 but no recommendation on a posology can be made.

Method of administration

Tablets:

Voriconazole film-coated tablets are to be taken at least one hour before, or one hour following, a meal.

Powder for solution for infusion:

Voriconazole requires reconstitution and dilution prior to administration as an intravenous infusion. Not for bolus injection.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 2.

Interacting drugs listed in this section and section 4.5 are a guide and not considered a comprehensive list of all possible drugs that may be contraindicated.

Coadministration of voriconazole is contraindicated with medicinal products that are highly dependent on CYP3A4 for metabolism, and for which elevated plasma concentrations are associated with serious and/or life-threatening reactions (see section 4.5):

- Terfenadine
- Astemizole
- Cisapride
- Pimozide
- Lurasidone
- Quinidine
- Ivabradine
- Ergot alkaloids (e.g., ergotamine, dihydroergotamine)
- Sirolimus
- Naloxegol
- Tolvaptan
- Finerenone
- Eplerenone
- Voclosporin
- Venetoclax: Coadministration contraindicated at initiation and during venetoclax dose titration phase.

Coadministration of voriconazole is contraindicated with medicinal products that induce CYP3A4 and significantly reduce voriconazole plasma concentrations:

- Coadministration with rifampicin, carbamazepine, long-acting barbiturates e.g., phenobarbital and St John's Wort (see section 4.5).

- Efavirenz:

Coadministration of standard doses of voriconazole with efavirenz doses of 400 mg once daily or higher is contraindicated(see section 4.5). For information on coadministration of voriconazole and lower doses of efavirenz see section 4.4.

- Ritonavir:

Coadministration with high-dose ritonavir (400 mg and above twice daily) is contraindicated (see section 4.5) For information on coadministration with lower doses of ritonavir see section 4.4.

4.4 Special warnings and precautions for use

Hypersensitivity

Caution should be used in prescribing voriconazole to patients with hypersensitivity to other azoles (see also section 4.8).

Powder for solution for infusion:

Duration of treatment

The duration of treatment with the intravenous formulation should be no longer than 6 months (see

section 6.1).

Cardiovascular

Voriconazole has been associated with QTc interval prolongation. There have been rare cases of torsades de pointes in patients taking voriconazole who had risk factors, such as history of cardiotoxic chemotherapy, cardiomyopathy, hypokalaemia and concomitant medicinal products that may have been contributory. Voriconazole should be administered with caution to patients with potentially proarrhythmic conditions, such as:

- Congenital or acquired QTc-prolongation.
- Cardiomyopathy, in particular when heart failure is present.
- Sinus bradycardia.
- Existing symptomatic arrhythmias.
- Concomitant medicinal product that is known to prolong QTc interval. Electrolyte disturbances such as hypokalaemia, hypomagnesaemia and hypocalcaemia should be monitored and corrected, if necessary, prior to initiation and during voriconazole therapy (see section 4.2). A study has been conducted in healthy volunteers which examined the effect on QTc interval of single doses of voriconazole up to 4 times the usual daily dose. No subject experienced an interval exceeding the potentially clinically relevant threshold of 500 msec (see section 5.1).

Powder for solution for infusion:

Infusion-related reactions

Infusion-related reactions, predominantly flushing and nausea have been observed during administration of the intravenous formulation of voriconazole. Depending on the severity of symptoms, consideration should be given to stopping treatment (see section 4.8).

Hepatic toxicity

In clinical trials, there have been cases of serious hepatic reactions during treatment with voriconazole (including clinical hepatitis, cholestasis and fulminant hepatic failure, including fatalities). Instances of hepatic reactions were noted to occur primarily in patients with serious underlying medical conditions (predominantly haematological malignancy). Transient hepatic reactions, including hepatitis and jaundice, have occurred among patients with no other identifiable risk factors. Liver dysfunction has usually been reversible on discontinuation of therapy (see section 4.8).

Monitoring of hepatic function

Patients receiving voriconazole must be carefully monitored for hepatic toxicity. Clinical management should include laboratory evaluation of hepatic function (specifically AST and ALT) at the initiation of treatment with voriconazole and at least weekly for the first month of treatment. Treatment duration should be as short as possible, however, if based on the benefit-risk assessment the treatment is continued (see section 4.2), monitoring frequency can be reduced to monthly if there are no changes in the liver function tests.

If the liver function tests become markedly elevated, voriconazole should be discontinued, unless the medical judgment of the risk-benefit of the treatment for the patient justifies continued use.

Monitoring of hepatic function should be carried out in both children and adults.

Serious dermatological adverse reactions

- Phototoxicity

In addition, voriconazole has been associated with phototoxicity including reactions such as ephelides, lentigo, actinic keratosis and pseudoporphyria. There is a potential increased risk of skin reactions/toxicity with concomitant use of photosensitising agents (e.g., methotrexate, etc). It is recommended that all patients, including children, avoid exposure to direct sunlight during voriconazole treatment and use measures such as protective clothing and sunscreen with high sun protection factor (SPF).

- Squamous cell carcinoma of the skin (SCC)

Squamous cell carcinoma of the skin (including cutaneous SCC in situ, or Bowen's disease) has been reported in patients, some of whom have reported prior phototoxic reactions. If phototoxic reactions occur multidisciplinary advice should be sought, voriconazole discontinuation and use of alternative antifungal agents should be considered and the patient should be referred to a dermatologist. If voriconazole is continued, however, dermatologic evaluation should be performed on a systematic and regular basis, to allow early detection and management of premalignant lesions. Voriconazole should be discontinued if premalignant skin lesions or squamous cell carcinoma are identified (see below the section under Long-term treatment).

- Severe cutaneous adverse reactions

Severe cutaneous adverse reactions (SCARs) including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug reaction with eosinophilia and systemic symptoms (DRESS), which can be life-threatening or fatal, have been reported with the use of voriconazole. If a patient develops a rash, he should be monitored closely and voriconazole discontinued if lesions progress.

Adrenal events

Reversible cases of adrenal insufficiency have been reported in patients receiving azoles including voriconazole. Adrenal insufficiency has been reported in patients receiving azoles with or without concomitant corticosteroids. In patients receiving azoles without corticosteroids, adrenal insufficiency is related to direct inhibition of steroidogenesis by azoles. In patients taking corticosteroids, voriconazole associated CYP3A4 inhibition of their metabolism may lead to corticosteroid excess and adrenal suppression (see section 4.5). Cushing's syndrome with and without subsequent adrenal insufficiency has also been reported in patients receiving voriconazole concomitantly with corticosteroids.

Patients on long-term treatment with voriconazole and corticosteroids (including inhaled corticosteroids e.g., budesonide and intranasal corticosteroids) should be carefully monitored for

adrenal cortex dysfunction both during treatment and when voriconazole is discontinued (see section 4.5). Patients should be instructed to seek immediate medical care if they develop signs and symptoms of Cushing's syndrome or adrenal insufficiency.

Long-term treatment

Long term exposure (treatment) greater than 180 days (6 months) requires careful assessment of the benefit-risk balance and physicians should therefore consider the need to limit the exposure to voriconazole (see sections 4.2 and 5.1).

Squamous cell carcinoma of the skin (SCC) (including cutaneous SCC in situ, or Bowen's disease) has been reported in relation with long-term voriconazole treatment (see section 4.8).

Non-infectious periostitis with elevated fluoride and alkaline phosphatase levels has been reported in transplant patients. If a patient develops skeletal pain and radiologic findings compatible with periostitis voriconazole discontinuation should be considered after multidisciplinary advice (see section 4.8).

Visual adverse reactions

There have been reports of prolonged visual adverse reactions, including blurred vision, optic neuritis and papilloedema (see section 4.8).

Renal adverse reactions

Acute renal failure has been observed in severely ill patients undergoing treatment with voriconazole. Patients being treated with voriconazole are likely to be treated concomitantly with nephrotoxic medicinal products and have concurrent conditions that may result in decreased renal function (see section 4.8).

Monitoring of renal function

Patients should be monitored for the development of abnormal renal function. This should include laboratory evaluation, particularly serum creatinine.

Monitoring of pancreatic function

Patients, especially children with risk factors for acute pancreatitis (e.g., recent chemotherapy, haematopoietic stem cell transplantation [HSCT]), should be monitored closely during voriconazole treatment. Monitoring of serum amylase or lipase may be considered in this clinical situation.

Paediatric population

Safety and effectiveness in paediatric subjects below the age of two years has not been established (see sections 4.8 and 5.1). Voriconazole is indicated for paediatric patients aged two years or older. A higher frequency of liver enzyme elevations was observed in the paediatric population (see section 4.8). Hepatic function should be monitored in both children and adults. Oral bioavailability

may be limited in paediatric patients aged 2 to <12 years with malabsorption and very low body weight for age. In that case, intravenous voriconazole administration is recommended.

- Serious dermatological adverse reactions (including SCC)

The frequency of phototoxicity reactions is higher in the paediatric population. As an evolution towards SCC has been reported, stringent measures for the photoprotection are warranted in this population of patients. In children experiencing photoaging injuries such as lentigines or ephelides, sun avoidance and dermatologic follow-up are recommended even after treatment discontinuation.

Phenytoin (CYP2C9 substrate and potent CYP450 inducer)

Careful monitoring of phenytoin levels is recommended when phenytoin is coadministered with voriconazole. Concomitant use of voriconazole and phenytoin should be avoided unless the benefit outweighs the risk (see section 4.5).

Efavirenz (CYP450 inducer; CYP3A4 inhibitor and substrate)

When voriconazole is coadministered with efavirenz the dose of voriconazole should be increased to 400 mg every 12 hours and the dose of efavirenz should be decreased to 300 mg every 24 hours (see sections 4.2, 4.3 and 4.5).

Glasdegib (CYP3A4 substrate)

Coadministration of voriconazole is expected to increase glasdegib plasma concentrations and increase the risk of QTc prolongation (see section 4.5). If concomitant use cannot be avoided, frequent ECG monitoring is recommended.

Tyrosine kinase inhibitors (CYP3A4 substrate)

Coadministration of voriconazole with tyrosine kinase inhibitors metabolised by CYP3A4 is expected to increase tyrosine kinase inhibitor plasma concentrations and the risk of adverse reactions. If concomitant use cannot be avoided, dose reduction of the tyrosine kinase inhibitor and close clinical monitoring is recommended (see section 4.5).

Rifabutin (potent CYP450 inducer)

Careful monitoring of full blood counts and adverse reactions to rifabutin (e.g., uveitis) is recommended when rifabutin is coadministered with voriconazole. Concomitant use of voriconazole and rifabutin should be avoided unless the benefit outweighs the risk (see section 4.5).

Ritonavir (potent CYP450 inducer, CYP3A4 inhibitor and substrate)

Coadministration of voriconazole and low-dose ritonavir (100 mg twice daily) should be avoided unless an assessment of the benefit/risk to the patient justifies the use of voriconazole (see sections 4.3 and 4.5).

Everolimus (CYP3A4 substrate, P-gp substrate)

Coadministration of voriconazole with everolimus is not recommended because voriconazole is expected to significantly increase everolimus concentrations. Currently there are insufficient data to allow dosing recommendations in this situation (see section 4.5).

Methadone (CYP3A4 substrate)

Frequent monitoring for adverse reactions and toxicity related to methadone, including QTc prolongation, is recommended when coadministered with voriconazole since methadone levels increased following coadministration of voriconazole. Dose reduction of methadone may be needed (see section 4.5).

Short-acting opiates (CYP3A4 substrate)

Reduction in the dose of alfentanil, fentanyl and other short acting opiates similar in structure to alfentanil and metabolised by CYP3A4 (e.g., sufentanil) should be considered when coadministered with voriconazole (see section 4.5). As the half-life of alfentanil is prolonged in a 4-fold manner when alfentanil is coadministered with voriconazole, and in an independent published study, concomitant use of voriconazole with fentanyl resulted in an increase in the mean $AUC_{0-\infty}$ of fentanyl, frequent monitoring for opiate-associated adverse reactions (including a longer respiratory monitoring period) may be necessary.

Long-acting opiates (CYP3A4 substrate)

Reduction in the dose of oxycodone and other long-acting opiates metabolised by CYP3A4 (e.g., hydrocodone) should be considered when coadministered with voriconazole. Frequent monitoring for opiate-associated adverse reactions may be necessary (see section 4.5).

Fluconazole (CYP2C9, CYP2C19 and CYP3A4 inhibitor)

Coadministration of oral voriconazole and oral fluconazole resulted in a significant increase in C_{max} and AUC_{τ} of voriconazole in healthy subjects. The reduced dose and/or frequency of voriconazole and fluconazole that would eliminate this effect have not been established. Monitoring for voriconazole associated adverse reactions is recommended if voriconazole is used sequentially after fluconazole (see section 4.5).

Excipients

Lactose

Tablet:

This medicinal product contains lactose and should not be given to patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption.

Sodium

This medicinal product contains less than 1 mmol sodium (23 mg) per tablet. Patients on low sodium diets should be informed that this medicinal product is essentially 'sodium-free'.

Powder for solution for infusion:

Sodium

This medicinal product contains 221 mg of sodium per vial, equivalent to 11% of the WHO recommended maximum daily intake of 2 g sodium for an adult.

Cyclodextrins

The powder for solution for infusion contains cyclodextrins (3,200 mg cyclodextrins in each vial which is equivalent to 160 mg/ml when reconstituted in 20 ml, see sections 2 and 6.1) which can influence the properties (such as toxicity) of the active substance and other medicines. Safety aspects of cyclodextrins have been considered during the development and safety assessment of the drug product.

As cyclodextrins are renally excreted, in patients with moderate to severe renal dysfunction accumulation of cyclodextrin may occur.

4.5 Drugs interactions

Voriconazole is metabolised by, and inhibits the activity of, cytochrome P450 isoenzymes, CYP2C19, CYP2C9, and CYP3A4. Inhibitors or inducers of these isoenzymes may increase or decrease voriconazole plasma concentrations, respectively, and there is potential for voriconazole to increase the plasma concentrations of substances metabolised by these CYP450 isoenzymes, in particular for substances metabolised by CYP3A4 since voriconazole is a strong CYP3A4 inhibitor though the increase in AUC is substrate dependent (see Table below).

Unless otherwise specified, drug interaction studies have been performed in healthy adult male subjects using multiple dosing to steady-state with oral voriconazole at 200 mg twice daily (BID). These results are relevant to other populations and routes of administration.

Voriconazole should be administered with caution in patients with concomitant medication that is known to prolong QTc interval. When there is also a potential for voriconazole to increase the plasma concentrations of substances metabolised by CYP3A4 isoenzymes (certain antihistamines, quinidine, cisapride, pimozide and ivabradine), coadministration is contraindicated (see below and section 4.3).

Interaction table

Interactions between voriconazole and other medicinal products are listed in the table below (once daily as “QD”, twice daily as “BID”, three times daily as “TID” and not determined as “ND”) ordered by therapeutic class. The direction of the arrow for each pharmacokinetic parameter is based on the 90% confidence interval of the geometric mean ratio being within (\leftrightarrow), below (\downarrow) or above (\uparrow) the 80-125% range. The asterisk (*) indicates a two-way interaction. AUC_{τ} , AUC_t and $AUC_{0-\infty}$ represent area under the curve over a dosing interval, from time zero to the time with detectable measurement and from time zero to infinity, respectively.

Medicinal products listed in the table are a guide and not considered a comprehensive list of all possible medicinal products that are contraindicated or may interact with voriconazole.

Medicinal product	Interaction geometric mean changes (%)	Recommendations concerning coadministration
Antacids		
Cimetidine (400 mg BID) <i>[non-specific CYP450 inhibitor and increases gastric pH]</i>	Voriconazole C _{max} ↑ 18% Voriconazole AUC _τ ↑ 23%	No dose adjustment
Omeprazole (40 mg QD)* <i>[CYP2C19 inhibitor; CYP2C19 and CYP3A4 substrate]</i>	Omeprazole C _{max} ↑ 116% Omeprazole AUC _τ ↑ 280% Voriconazole C _{max} ↑ 15% Voriconazole AUC _τ ↑ 41% Other proton pump inhibitors that are CYP2C19 substrates may also be inhibited by voriconazole and may result in increased plasma concentrations of these medicinal products.	No dose adjustment of voriconazole is recommended. When initiating voriconazole in patients already receiving omeprazole doses of 40 mg or above, it is recommended that the omeprazole dose be halved.
Ranitidine (150 mg BID) <i>[increases gastric pH]</i>	Voriconazole C _{max} and AUC _τ ↔	No dose adjustment
Antiarrhythmics		
Digoxin (0.25 mg QD) <i>[P-gp substrate]</i>	Digoxin C _{max} ↔ Digoxin AUC _τ ↔	No dose adjustment
Quinidine <i>[CYP3A4 substrate]</i>	Although not studied, increased plasma concentrations of quinidine can lead to QTc prolongation and rare occurrences of torsades de pointes.	Contraindicated (see section 4.3)
Antibacterials		
Flucloxacillin <i>[CYP450 inducer]</i>	Significantly decreased plasma voriconazole concentrations have been reported.	If concomitant administration of voriconazole with flucloxacillin cannot be avoided, monitor for potential loss of voriconazole effectiveness (e.g., by therapeutic drug monitoring); increasing the dose of voriconazole may be needed.
Macrolide antibiotics Azithromycin (500 mg QD) Erythromycin (1 g BID) <i>[CYP3A4 inhibitor]</i>	Voriconazole C _{max} and AUC _τ ↔ Voriconazole C _{max} and AUC _τ ↔ The effect of voriconazole on either erythromycin or azithromycin is unknown.	No dose adjustment

Rifabutin <i>[potent CYP450 inducer]</i> 300 mg QD 300 mg QD (coadministered with voriconazole 350 mg BID)* 300 mg QD (coadministered with voriconazole 400 mg BID)*	Voriconazole C _{max} ↓ 69% Voriconazole AUC _τ ↓ 78% Compared to voriconazole 200 mg BID, Voriconazole C _{max} ↓ 4% Voriconazole AUC _τ ↓ 32% Rifabutin C _{max} ↑ 195% Rifabutin AUC _τ ↑ 331% Compared to voriconazole 200 mg BID, Voriconazole C _{max} ↑ 104% Voriconazole AUC _τ ↑ 87%	Concomitant use of voriconazole and rifabutin should be avoided unless the benefit outweighs the risk. The maintenance dose of voriconazole may be increased to 5 mg/kg intravenously BID or from 200 mg to 350 mg orally BID (100 mg to 200 mg orally BID in patients less than 40 kg) (see section 4.2). Careful monitoring of full blood counts and adverse reactions to rifabutin (e.g., uveitis) is recommended when rifabutin is coadministered with voriconazole.
Rifampicin (600 mg QD) <i>[potent CYP450 inducer]</i>	Voriconazole C _{max} ↓ 93% Voriconazole AUC _τ ↓ 96%	Contraindicated (see section 4.3)
Anti-cancer agents		
Glasdegib <i>[CYP3A4 substrate]</i>	Although not studied, voriconazole is likely to increase the plasma concentrations of glasdegib and increase risk of QTc prolongation.	If concomitant use cannot be avoided, frequent ECG monitoring is recommended (see section 4.4).
Tretinoin <i>[CYP3A4 substrate]</i>	Although not studied, voriconazole may increase tretinoin concentrations and increase risk of adverse reactions (pseudotumor cerebri, hypercalcaemia).	Dose adjustment of tretinoin is recommended during treatment with voriconazole and after its discontinuation.
Tyrosine kinase inhibitors (including but not limited to: axitinib, bosutinib, cabozantinib, ceritinib, cobimetinib, dabrafenib, dasatinib, nilotinib, sunitinib, ibrutinib, ribociclib) <i>[CYP3A4 substrates]</i>	Although not studied, voriconazole may increase plasma concentrations of tyrosine kinase inhibitors metabolised by CYP3A4.	If concomitant use cannot be avoided, dose reduction of the tyrosine kinase inhibitor and close clinical monitoring is recommended (see section 4.4).
Venetoclax <i>[CYP3A substrate]</i>	Although not studied, voriconazole is likely to significantly increase the plasma concentrations of venetoclax.	Concomitant administration of voriconazole is contraindicated at initiation and during venetoclax dose titration phase (see section 4.3). Dose reduction of venetoclax is required as instructed in venetoclax prescribing information during steady daily dosing; close monitoring for signs of toxicity is recommended.

Vinca Alkaloids (including but not limited to: vincristine and vinblastine) [CYP3A4 substrates]	Although not studied, voriconazole is likely to increase the plasma concentrations of vinca alkaloids and lead to neurotoxicity.	Dose reduction of vinca alkaloids should be considered.
Anticoagulants		
Warfarin (30 mg single dose, coadministered with 300 mg BID voriconazole) [CYP2C9 substrate]	Maximum increase in prothrombin time was approximately 2-fold.	Close monitoring of prothrombin time or other suitable anticoagulation tests is recommended, and the dose of anticoagulants should be adjusted accordingly.
Other oral coumarins (including but not limited to: phenprocoumon, acenocoumarol) [CYP2C9 and CYP3A4 substrates]	Although not studied, voriconazole may increase the plasma concentrations of coumarins that may cause an increase in prothrombin time.	
Anticonvulsants		
Carbamazepine and long-acting barbiturates (including but not limited to: phenobarbital, mephobarbital) [potent CYP450 inducers]	Although not studied, carbamazepine and long-acting barbiturates are likely to significantly decrease plasma voriconazole concentrations.	Contraindicated (see section 4.3)
Phenytoin [CYP2C9 substrate and potent CYP450 inducer]		Concomitant use of voriconazole and phenytoin should be avoided unless the benefit outweighs the risk. Careful monitoring of phenytoin plasma levels is recommended.
300 mg QD	Voriconazole C _{max} ↓ 49% Voriconazole AUC _τ ↓ 69%	
300 mg QD (coadministered with voriconazole 400 mg BID)*	Phenytoin C _{max} ↑ 67% Phenytoin AUC _τ ↑ 81% Compared to voriconazole 200 mg BID, Voriconazole C _{max} ↑ 34% Voriconazole AUC _τ ↑ 39%	Phenytoin may be coadministered with voriconazole if the maintenance dose of voriconazole is increased to 5 mg/kg IV BID or from 200 mg to 400 mg oral BID (100 mg to 200 mg oral BID in patients less than 40 kg) (see section 4.2).
Antidiabetics		
Sulfonylureas (including but not limited to: tolbutamide, glipizide, glyburide) [CYP2C9 substrates]	Although not studied, voriconazole is likely to increase the plasma concentrations of sulfonylureas and cause hypoglycaemia.	Careful monitoring of blood glucose is recommended. Dose reduction of sulfonylureas should be considered.

Anti-fungals		
Fluconazole (200 mg QD) [CYP2C9, CYP2C19 and CYP3A4 inhibitor]	Voriconazole C _{max} ↑ 57% Voriconazole AUC _τ ↑ 79% Fluconazole C _{max} ND Fluconazole AUC _τ ND	The reduced dose and/or frequency of voriconazole and fluconazole that would eliminate this effect have not been established. Monitoring for voriconazole-associated adverse reactions is recommended if voriconazole is used sequentially after fluconazole.
Antihistamines		
Astemizole [CYP3A4 substrate]	Although not studied, increased plasma concentrations of astemizole can lead to QTc prolongation and rare occurrences of torsades de pointes.	Contraindicated (see section 4.3)
Terfenadine [CYP3A4 substrate]	Although not studied, increased plasma concentrations of terfenadine can lead to QTc prolongation and rare occurrences of torsades de pointes.	Contraindicated (see section 4.3)
Anti HIV agents		
Indinavir (800 mg TID) [CYP3A4 inhibitor and substrate]	Indinavir C _{max} ↔ Indinavir AUC _τ ↔ Voriconazole C _{max} ↔ Voriconazole AUC _τ ↔	No dose adjustment
Ritonavir (protease inhibitor) [potent CYP450 inducer; CYP3A4 inhibitor and substrate]		
High dose (400 mg BID)	Ritonavir C _{max} and AUC _τ ↔ Voriconazole C _{max} ↓ 66% Voriconazole AUC _τ ↓ 82%	Coadministration of voriconazole and high doses of ritonavir (400 mg and above BID) is contraindicated (see section 4.3).
Low dose (100 mg BID)*	Ritonavir C _{max} ↓ 25% Ritonavir AUC _τ ↓ 13% Voriconazole C _{max} ↓ 24% Voriconazole AUC _τ ↓ 39%	Coadministration of voriconazole and low-dose ritonavir (100 mg BID) should be avoided unless an assessment of the benefit/risk to the patient justifies the use of voriconazole.
Other HIV Protease Inhibitors (including but not limited to: saquinavir, amprenavir and nelfinavir)* [CYP3A4 substrates and inhibitors]	Not studied clinically. <i>In vitro</i> studies show that voriconazole may inhibit the metabolism of HIV protease inhibitors and the metabolism of voriconazole may also be inhibited by HIV protease inhibitors.	Careful monitoring for any occurrence of drug toxicity and/or lack of efficacy, and dose adjustment may be needed.

<p>Efavirenz (a non-nucleoside reverse transcriptase inhibitor, (NNRTI)) [<i>CYP450 inducer; CYP3A4 inhibitor and substrate</i>]</p> <p>Efavirenz 400 mg QD, coadministered with voriconazole 200 mg BID*</p> <p>Efavirenz 300 mg QD, coadministered with voriconazole 400 mg BID*</p>	<p>Efavirenz C_{max} ↑ 38% Efavirenz AUC_{τ} ↑ 44% Voriconazole C_{max} ↓ 61% Voriconazole AUC_{τ} ↓ 77%</p> <p>Compared to efavirenz 600 mg QD, Efavirenz C_{max} ↔ Efavirenz AUC_{τ} ↑ 17%</p> <p>Compared to voriconazole 200 mg BID, Voriconazole C_{max} ↑ 23% Voriconazole AUC_{τ} ↓ 7%</p>	<p>Use of standard doses of voriconazole with efavirenz doses of 400 mg QD or higher is contraindicated (see section 4.3).</p> <p>Voriconazole may be coadministered with efavirenz if the voriconazole maintenance dose is increased to 400 mg BID and the efavirenz dose is decreased to 300 mg QD. When voriconazole treatment is stopped, the initial dose of efavirenz should be restored (see sections 4.2 and 4.4).</p>
<p>Other Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs) (including but not limited to: delavirdine, nevirapine)* [<i>CYP3A4 substrates, inhibitors or CYP450 inducers</i>]</p>	<p>Not studied clinically. <i>In vitro</i> studies show that the metabolism of voriconazole may be inhibited by NNRTIs and voriconazole may inhibit the metabolism of NNRTIs. The findings of the effect of efavirenz on voriconazole suggest that the metabolism of voriconazole may be induced by an NNRTI.</p>	<p>Careful monitoring for any occurrence of drug toxicity and/or lack of efficacy, and dose adjustment may be needed.</p>
<p>Antipsychotics</p>		
<p>Lurasidone [<i>CYP3A4 substrate</i>]</p>	<p>Although not studied, voriconazole is likely to significantly increase the plasma concentrations of lurasidone.</p>	<p>Contraindicated (see section 4.3)</p>
<p>Pimozide [<i>CYP3A4 substrate</i>]</p>	<p>Although not studied, increased plasma concentrations of pimozide can lead to QTc prolongation and rare occurrences of torsades de pointes.</p>	<p>Contraindicated (see section 4.3)</p>
<p>Anti virals</p>		
<p>Letemovir [<i>CYP2C9 and CYP2C19 inducer</i>]</p>	<p>Voriconazole C_{max} ↓ 39% Voriconazole AUC_{0-12} ↓ 44% Voriconazole C_{12} ↓ 51%</p>	<p>If concomitant administration of voriconazole with letemovir cannot be avoided, monitor for loss of voriconazole effectiveness.</p>

<i>Benzodiazepines</i>		
<p>[CYP3A4 substrates] Midazolam (0.05 mg/kg IV single dose)</p>	<p>In an independent published study, Midazolam AUC_{0-∞} ↑ 3.7-fold</p>	<p>Dose reduction of benzodiazepines should be considered.</p>
<p>Midazolam (7.5 mg oral single dose)</p>	<p>In an independent published study, Midazolam C_{max} ↑ 3.8-fold Midazolam AUC_{0-∞} ↑ 10.3-fold</p>	
<p>Other benzodiazepines (including but not limited to: triazolam, alprazolam)</p>	<p>Although not studied, voriconazole is likely to increase the plasma concentrations of other benzodiazepines that are metabolised by CYP3A4 and lead to a prolonged sedative effect.</p>	
<i>Cardiovascular agents</i>		
<p>Ivabradine [CYP3A4 substrates]</p>	<p>Although not studied, increased plasma concentrations of ivabradine can lead to QTc prolongation and rare occurrences of torsades de pointes.</p>	<p>Contraindicated (see section 4.3)</p>
<i>Cystic fibrosis transmembrane conductance regulator potentiators</i>		
<p>Ivacaftor [CYP3A4 substrate]</p>	<p>Although not studied, voriconazole is likely to increase the plasma concentrations of ivacaftor with risk of increased adverse reactions.</p>	<p>Dose reduction of ivacaftor is recommended.</p>
<i>Ergot derivatives</i>		
<p>Ergot alkaloids (including but not limited to: ergotamine and dihydroergotamine) [CYP3A4 substrates]</p>	<p>Although not studied, voriconazole is likely to increase the plasma concentrations of ergot alkaloids and lead to ergotism.</p>	<p>Contraindicated (see section 4.3)</p>
<i>GI motility agents</i>		
<p>Cisapride [CYP3A4 substrate]</p>	<p>Although not studied, increased plasma concentrations of cisapride can lead to QTc prolongation and rare occurrences of torsades de pointes.</p>	<p>Contraindicated (see section 4.3)</p>
<i>Herbal medicines</i>		
<p>St. John's Wort [CYP450 inducer; P-gp inducer] 300 mg TID (coadministered with voriconazole 400 mg single dose)</p>	<p>In an independent published study, Voriconazole AUC_{0-∞} ↓ 59%</p>	<p>Contraindicated (see section 4.3)</p>

Immunosuppressants		
<i>[CYP3A4 substrates]</i>		
Ciclosporin (in stable renal transplant recipients receiving chronic ciclosporin therapy)	Ciclosporin C_{max} ↑ 13% Ciclosporin AUC_{τ} ↑ 70%	When initiating voriconazole in patients already on ciclosporin it is recommended that the ciclosporin dose be halved and ciclosporin level carefully monitored. Increased ciclosporin levels have been associated with nephrotoxicity. <u>When voriconazole is discontinued, ciclosporin levels must be carefully monitored and the dose increased as necessary.</u>
Everolimus <i>[also P-gp substrate]</i>	Although not studied, voriconazole is likely to significantly increase the plasma concentrations of everolimus.	Coadministration of voriconazole and everolimus is not recommended because voriconazole is expected to significantly increase everolimus concentrations (see section 4.4).
Sirolimus (2 mg single dose)	In an independent published study, Sirolimus C_{max} ↑ 6.6-fold Sirolimus $AUC_{0-\infty}$ ↑ 11-fold	Coadministration of voriconazole and sirolimus is contraindicated (see section 4.3).
Tacrolimus (0.1 mg/kg single dose)	Tacrolimus C_{max} ↑ 117% Tacrolimus AUC_{τ} ↑ 221%	When initiating voriconazole in patients already on tacrolimus, it is recommended that the tacrolimus dose be reduced to a third of the original dose and tacrolimus level carefully monitored. Increased tacrolimus levels have been associated with nephrotoxicity. <u>When voriconazole is discontinued, tacrolimus levels must be carefully monitored and the dose increased as necessary.</u>
Voclosporin	Although not studied, voriconazole is likely to significantly increase the plasma concentrations of voclosporin.	Contraindicated (see section 4.3)
Mycophenolic acid (1 g single dose) <i>[UDP-glucuronyl transferase substrate]</i>	Mycophenolic acid C_{max} ↔ Mycophenolic acid AUC_{τ} ↔	No dose adjustment
Lipid lowering agents/HMG- CoA reductase inhibitors		

Statins (e.g., lovastatin) [CYP3A4 substrates]	Although not studied, voriconazole is likely to increase the plasma concentrations of statins that are metabolised by CYP3A4 and could lead to rhabdomyolysis.	If concomitant administration of voriconazole with statins metabolised by CYP3A4 cannot be avoided, dose reduction of the statin should be considered.
<i>Non-steroidal selective mineralocorticoid receptor (MR) antagonists</i>		
Finerenone [CYP3A4 substrate]	Although not studied, voriconazole is likely to significantly increase the plasma concentrations of finerenone.	Contraindicated (see section 4.3)
Eplerenone [CYP3A4 substrate]	Although not studied, voriconazole is likely to significantly increase the plasma concentrations of eplerenone.	Contraindicated (see section 4.3)
<i>Non-steroidal anti-inflammatory drugs (NSAIDs)</i>		
[CYP2C9 substrates] Ibuprofen (400 mg single dose) Diclofenac (50 mg single dose)	S-Ibuprofen C _{max} ↑ 20% S-Ibuprofen AUC _{0-∞} ↑ 100% Diclofenac C _{max} ↑ 114% Diclofenac AUC _{0-∞} ↑ 78%	Frequent monitoring for adverse reactions and toxicity related to NSAIDs is recommended. Dose reduction of NSAIDs may be needed.
<i>Opioids</i>		
Long-Acting Opiates [CYP3A4 substrates] Oxycodone (10 mg single dose)	In an independent published study, Oxycodone C _{max} ↑ 1.7-fold Oxycodone AUC _{0-∞} ↑ 3.6-fold	Dose reduction in oxycodone and other long-acting opiates metabolised by CYP3A4 (e.g., hydrocodone) should be considered. Frequent monitoring for opiate-associated adverse reactions may be necessary.
Methadone (32-100 mg QD) [CYP3A4 substrate]	R-methadone (active) C _{max} ↑ 31% R-methadone (active) AUC _τ ↑ 47% S-methadone C _{max} ↑ 65% S-methadone AUC _τ ↑ 103%	Frequent monitoring for adverse reactions and toxicity related to methadone, including QTc prolongation, is recommended. Dose reduction of methadone may be needed.
Short-acting Opiates [CYP3A4 substrates] Alfentanil (20 µg/kg single dose, with concomitant naloxone) Fentanyl (5 µg/kg single dose)	In an independent published study, Alfentanil AUC _{0-∞} ↑ 6-fold In an independent published study, Fentanyl AUC _{0-∞} ↑ 1.34-fold	Dose reduction of alfentanil, fentanyl and other short-acting opiates similar in structure to alfentanil and metabolised by CYP3A4 (e.g., sufentanil) should be considered. Extended and frequent monitoring for respiratory depression and other opiate-associated adverse reactions is recommended.
<i>Opioid receptor antagonists</i>		
Naloxegol [CYP3A4 substrate]	Although not studied, voriconazole is likely to significantly increase the plasma concentrations of naloxegol.	Contraindicated (see section 4.3)

Oral contraceptives		
Oral Contraceptives* [CYP3A4 substrate; CYP2C19 inhibitor] Norethisterone/ethinylestradiol (1 mg/0.035 mg QD)	Ethinylestradiol C _{max} ↑ 36% Ethinylestradiol AUC _τ ↑ 61% Norethisterone C _{max} ↑ 15% Norethisterone AUC _τ ↑ 53% Voriconazole C _{max} ↑ 14% Voriconazole AUC _τ ↑ 46%	Monitoring for adverse reactions related to oral contraceptives, in addition to those for voriconazole, is recommended.
Steroids		
Corticosteroids Prednisolone (60 mg single dose) [CYP3A4 substrate]	Prednisolone C _{max} ↑ 11% Prednisolone AUC _{0-∞} ↑ 34%	No dose adjustment Patients on long-term treatment with voriconazole and corticosteroids (including inhaled corticosteroids e.g., budesonide and intranasal corticosteroids) should be carefully monitored for adrenal cortex dysfunction both during treatment and when voriconazole is discontinued (see section 4.4).
Vasopressin receptor antagonists		
Tolvaptan [CYP3A substrate]	Although not studied, voriconazole is likely to significantly increase the plasma concentrations of tolvaptan.	Contraindicated (see section 4.3)

4.6 Use in special populations

Pregnancy

There are no adequate data on the use of voriconazole in pregnant women available.

Studies in animals have shown reproductive toxicity (see section 6.1). The potential risk for humans is unknown.

Voriconazole must not be used during pregnancy unless the benefit to the mother clearly outweighs the potential risk to the foetus.

Women of child-bearing potential

Women of child-bearing potential must always use effective contraception during treatment.

Breast-feeding

The excretion of voriconazole into breast milk has not been investigated. Breast-feeding must be stopped on initiation of treatment with voriconazole.

Fertility

In an animal study, no impairment of fertility was demonstrated in male and female rats (see section 6.1).

4.7 Effects on ability to drive and use machines

Voriconazole has moderate influence on the ability to drive and use machines. It may cause transient and reversible changes to vision, including blurring, altered/enhanced visual perception, and/or photophobia. Patients must avoid potentially hazardous tasks, such as driving or operating machinery while experiencing these symptoms.

4.8 Undesirable effects

Summary of safety profile

The safety profile of voriconazole in adults is based on an integrated safety database of more than 2,000 subjects (including 1,603 adult patients in therapeutic trials) and an additional 270 adults in prophylaxis trials. This represents a heterogeneous population, containing patients with haematological malignancy; HIV infected patients with oesophageal candidiasis and refractory fungal infections, non-neutropenic patients with candidaemia or aspergillosis and healthy volunteers.

The most commonly reported adverse reactions were visual impairment, pyrexia, rash, vomiting, nausea, diarrhoea, headache, peripheral oedema, liver function test abnormal, respiratory distress and abdominal pain

The severity of the adverse reactions was generally mild to moderate. No clinically significant differences were seen when the safety data were analysed by age, race, or gender.

Tabulated list of adverse reactions

In the table below, since the majority of the studies were of an open nature, all causality adverse reactions and their frequency categories in 1,873 adults from pooled therapeutic (1,603) and prophylaxis (270) studies, by system organ class, are listed.

Frequency categories are expressed as: Very common ($\geq 1/10$); Common ($\geq 1/100$ to $< 1/10$); Uncommon ($\geq 1/1,000$ to $< 1/100$); Rare ($\geq 1/10,000$ to $< 1/1,000$); Very rare ($< 1/10,000$); Not known (cannot be estimated from the available data).

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Undesirable effects reported in subjects receiving voriconazole:

System Organ Class	Very common ≥1/10	Common ≥1/100 to <1/10	Uncommon ≥1/1,000 to <1/100	Rare ≥1/10,000 to <1/1,000	Frequency not known (cannot be estimated from available data)
Infections and infestations		sinusitis	pseudomembranous colitis		
Neoplasms benign, malignant and unspecified (including cysts and polyps)		squamous cell carcinoma (including cutaneous SCC in situ, or Bowen's disease)*,**			
Blood and lymphatic system disorders		agranulocytosis ¹ , pancytopenia, thrombocytopenia ² , leukopenia, anaemia	bone marrow failure, lymphadenopathy, eosinophilia	disseminated intravascular coagulation	
Immune system disorders			hypersensitivity	anaphylactoid reaction	
Endocrine disorders			adrenal insufficiency, hypothyroidism	hyperthyroidism	
Metabolism and nutrition disorders	oedema peripheral	hypoglycaemia, hypokalaemia, hyponatraemia			
Psychiatric disorders		depression, hallucination, anxiety, insomnia, agitation, confusional state			
Nervous system disorders	Headache	convulsion, syncope, tremor, hypertonia ³ , paraesthesia, somnolence, dizziness	brain oedema, encephalopathy ⁴ , extrapyramidal disorder ⁵ , neuropathy peripheral, ataxia, hypoaesthesia, dysgeusia	hepatic encephalopathy, Guillain-Barre syndrome, nystagmus	

System Organ Class	Very common ≥1/10	Common ≥1/100 to <1/10	Uncommon ≥1/1,000 to <1/100	Rare ≥1/10,000 to <1/1,000	Frequency not known (cannot be estimated from available data)
Eye disorders	visual impairment ⁶	retinal haemorrhage	optic nerve disorder ⁷ , papilloedema ⁸ , oculogyric crisis, diplopia, scleritis, blepharitis	optic atrophy, corneal opacity	
Ear and labyrinth disorders			hyposacusis, vertigo, tinnitus		
Cardiac disorders		arrhythmia supraventricular, tachycardia, bradycardia	ventricular fibrillation, ventricular extrasystoles, ventricular tachycardia, electrocardiogram QT prolonged, supraventricular tachycardia	<i>torsades de pointes</i> , atrioventricular block complete, bundle branch block, nodal rhythm	
Vascular disorders		hypotension, phlebitis	thrombophlebitis, lymphangitis		
Respiratory, thoracic and mediastinal disorders	respiratory distress ⁹	acute respiratory distress syndrome, pulmonary oedema			
Gastrointestinal disorders	diarrhoea, vomiting, abdominal pain, nausea	cheilitis, dyspepsia, constipation, gingivitis	peritonitis, pancreatitis, swollen tongue, duodenitis, gastroenteritis, glossitis		
Hepatobiliary disorders	liver function test abnormal	jaundice, jaundice cholestatic, hepatitis ¹⁰	hepatic failure, hepatomegaly, cholecystitis, cholelithiasis		
Skin and subcutaneous tissue disorders	Rash	dermatitis exfoliative, alopecia, rash maculopapular, pruritus, erythema, phototoxicity**	Stevens-Johnson syndrome ⁸ , purpura, urticaria, dermatitis allergic, rash papular, rash macular, eczema	toxic epidermal necrolysis ⁸ , drug reaction with eosinophilia and systemic symptoms (DRESS) ⁸ , angioedema,	cutaneous lupus erythematosus*, ephelides*, lentigo*

System Organ Class	Very common ≥1/10	Common ≥1/100 to <1/10	Uncommon ≥1/1,000 to <1/100	Rare ≥1/10,000 to <1/1,000	Frequency not known (cannot be estimated from available data)
				actinic keratosis*, pseudoporphyria erythema multiforme, psoriasis, drug eruption	
Musculoskeletal and connective tissue disorders		back pain	arthritis, periostitis*,**		
Renal and urinary disorders		renal failure acute, haematuria	renal tubular necrosis, proteinuria, nephritis		
General disorders and administration site conditions	Pyrexia	chest pain, face oedema ¹¹ , asthenia, chills	infusion site reaction, influenza like illness		
Investigations		blood creatinine increased	blood urea increased, blood cholesterol increased		

*ADR identified post-marketing

**Frequency category is based on an observational study utilising real-world data from secondary data sources in Sweden

¹ Includes febrile neutropenia and neutropenia.

² Includes immune thrombocytopenic purpura.

³ Includes nuchal rigidity and tetany.

⁴ Includes hypoxic-ischaemic encephalopathy and metabolic encephalopathy.

⁵ Includes akathisia and parkinsonism.

⁶ See “Visual impairments” paragraph in section 4.8.

⁷ Prolonged optic neuritis has been reported post-marketing. See section 4.4.

⁸ See section 4.4.

⁹ Includes dyspnoea and dyspnoea exertional.

¹⁰ Includes drug-induced liver injury, hepatitis toxic, hepatocellular injury and hepatotoxicity.

¹¹ Includes periorbital oedema, lip oedema, and oedema mouth.

Description of selected adverse reactions

Visual impairments

In clinical trials, visual impairments (including blurred vision, photophobia, chloropsia, chromatopsia, colour blindness, cyanopsia, eye disorder, halo vision, night blindness, oscillopsia, photopsia, scintillating scotoma, visual acuity reduced, visual brightness, visual field defect,

vitreous floaters, and xanthopsia) with voriconazole were very common. These visual impairments were transient and fully reversible, with the majority spontaneously resolving within 60 minutes and no clinically significant long-term visual effects were observed. There was evidence of attenuation with repeated doses of voriconazole. The visual impairments were generally mild, rarely resulted in discontinuation and were not associated with long-term sequelae. Visual impairments may be associated with higher plasma concentrations and/or doses.

The mechanism of action is unknown, although the site of action is most likely to be within the retina. In a study in healthy volunteers investigating the impact of voriconazole on retinal function, voriconazole caused a decrease in the electroretinogram (ERG) waveform amplitude. The ERG measures electrical currents in the retina. The ERG changes did not progress over 29 days of treatment and were fully reversible on withdrawal of voriconazole.

There have been post-marketing reports of prolonged visual adverse events (see section 4.4).

Dermatological reactions

Dermatological reactions were very common in patients treated with voriconazole in clinical trials, but these patients had serious underlying diseases and were receiving multiple concomitant medicinal products. The majority of rashes were of mild to moderate severity. Patients have developed severe cutaneous adverse reactions (SCARs), including Stevens-Johnson syndrome (SJS) (uncommon), toxic epidermal necrolysis (TEN) (rare), drug reaction with eosinophilia and systemic symptoms (DRESS) (rare) and erythema multiforme (rare) during treatment with voriconazole (see section 4.4).

If a patient develops a rash they should be monitored closely and voriconazole discontinued if lesions progress. Photosensitivity reactions such as ephelides, lentigo and actinic keratosis have been reported, especially during long-term therapy (see section 4.4).

There have been reports of squamous cell carcinoma of the skin (including cutaneous SCC in situ, or Bowen's disease) in patients treated with voriconazole for long periods of time; the mechanism has not been established (see section 4.4).

Liver function tests

The overall incidence of transaminase increases $>3 \times \text{ULN}$ (not necessarily comprising an adverse event) in the voriconazole clinical programme was 18.0% (319/1,768) in adults and 25.8% (73/283) in paediatric subjects who received voriconazole for pooled therapeutic and prophylaxis use. Liver function test abnormalities may be associated with higher plasma concentrations and/or doses. The majority of abnormal liver function tests either resolved during treatment without dose adjustment or following dose adjustment, including discontinuation of therapy.

Voriconazole has been associated with cases of serious hepatic toxicity, in patients with other serious underlying conditions. This includes cases of jaundice, hepatitis and hepatic failure leading to death (see section 4.4).

Infusion-related reactions

During infusion of the intravenous formulation of voriconazole in healthy subjects, anaphylactoid-type reactions, including flushing, fever, sweating, tachycardia, chest tightness, dyspnea, faintness, nausea, pruritus and rash have occurred. Symptoms appeared immediately upon initiating the infusion (see section 4.4).

Paediatric population

The safety of voriconazole was investigated in 288 paediatric patients aged 2 to <12 years (169) and 12 to <18 years (119) who received voriconazole for prophylaxis (183) and therapeutic use (105) in clinical trials. The safety of voriconazole was also investigated in 158 additional paediatric patients aged 2 to <12 years in compassionate use programs. Overall, the safety profile of voriconazole in paediatric population was similar to that in adults. However, a trend towards a higher frequency of liver enzyme elevations, reported as adverse events in clinical trials was observed in paediatric patients as compared to adults (14.2% transaminases increased in paediatrics compared to 5.3% in adults). Post-marketing data suggest there might be a higher occurrence of skin reactions (especially erythema) in the paediatric population compared to adults. In the 22 patients less than 2 years old who received voriconazole in a compassionate use programme, the following adverse reactions (for which a relationship to voriconazole could not be excluded) were reported: photosensitivity reaction (1), arrhythmia (1), pancreatitis (1), blood bilirubin increased (1), hepatic enzymes increased (1), rash (1) and papilloedema (1). There have been post-marketing reports of pancreatitis in paediatric patients.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions.

4.9 Overdose

In clinical trials there were 3 cases of accidental overdose. All occurred in paediatric patients who received up to five times the recommended intravenous dose of voriconazole. A single adverse reaction of photophobia of 10 minutes duration was reported.

There is no known antidote to voriconazole.

Voriconazole is haemodialysed with a clearance of 121 ml/min. The intravenous vehicle, SBECD, is haemodialysed with a clearance of 55 ml/min. In an overdose, haemodialysis may assist in the removal of voriconazole and SBECD from the body.

5. PHARMACOLOGICAL PROPERTIES

5.1 Mechanism of action

Pharmacotherapeutic group: Antimycotics for systemic use, triazole derivatives, ATC code: J02A C03.

Voriconazole is a triazole antifungal agent. The primary mode of action of voriconazole is the inhibition of fungal cytochrome P-450-mediated 14 alpha-lanosterol demethylation, an essential step in fungal ergosterol biosynthesis. The accumulation of 14 alpha-methyl sterols correlates with the subsequent loss of ergosterol in the fungal cell membrane and may be responsible for the antifungal activity of voriconazole. Voriconazole has been shown to be more selective for fungal cytochrome P-450 enzymes than for various mammalian cytochrome P-450 enzyme systems.

5.2 Pharmacodynamic properties

Pharmacokinetic/Pharmacodynamic relationship

In 10 therapeutic studies, the median for the average and maximum plasma concentrations in individual subjects across the studies was 2425 ng/ml (inter-quartile range 1193 to 4380 ng/ml) and 3742 ng/ml (inter-quartile range 2027 to 6302 ng/ml), respectively. A positive association between mean, maximum or minimum plasma voriconazole concentration and efficacy in therapeutic studies was not found and this relationship has not been explored in prophylaxis studies.

Pharmacokinetic-Pharmacodynamic analyses of clinical trial data identified positive associations between plasma voriconazole concentrations and both liver function test abnormalities and visual disturbances. Dose adjustments in prophylaxis studies have not been explored.

Clinical efficacy and safety

In vitro, voriconazole displays broad-spectrum antifungal activity with antifungal potency against *Candida* species (including fluconazole resistant *C. krusei* and resistant strains of *C. glabrata* and *C. albicans*) and fungicidal activity against all *Aspergillus* species tested. In addition voriconazole shows *in vitro* fungicidal activity against emerging fungal pathogens, including those such as *Scedosporium* or *Fusarium* which have limited susceptibility to existing antifungal agents.

Clinical efficacy defined as partial or complete response, has been demonstrated for *Aspergillus* spp. including *A. flavus*, *A. fumigatus*, *A. terreus*, *A. niger*, *A. nidulans*, *Candida* spp., including *C. albicans*, *C. glabrata*, *C. krusei*, *C. parapsilosis* and *C. tropicalis* and limited numbers of *C. dubliniensis*, *C. inconspicua*, and *C. guilliermondii*, *Scedosporium* spp., including *S. apiospermum*, *S. prolificans* and *Fusarium* spp.

Other treated fungal infections (often with either partial or complete response) included isolated cases of *Alternaria* spp., *Blastomyces dermatitidis*, *Blastoschizomyces capitatus*, *Cladosporium* spp., *Coccidioides immitis*, *Conidiobolus coronatus*, *Cryptococcus neoformans*, *Exserohilum rostratum*, *Exophiala spinifera*, *Fonsecaea pedrosoi*, *Madurella mycetomatis*, *Paecilomyces lilacinus*, *Penicillium* spp. including *P. marneffeii*, *Phialophora richardsiae*, *Scopulariopsis brevicaulis* and *Trichosporon* spp. including *T. beigelii* infections.

In vitro activity against clinical isolates has been observed for *Acremonium* spp., *Alternaria* spp., *Bipolaris* spp., *Cladophialophora* spp., and *Histoplasma capsulatum*, with most strains being inhibited by concentrations of voriconazole in the range 0.05 to 2 µg/ml.

In vitro activity against the following pathogens has been shown, but the clinical significance is unknown: *Curvularia* spp. and *Sporothrix* spp.

Breakpoints

Specimens for fungal culture and other relevant laboratory studies (serology, histopathology) should be obtained prior to therapy to isolate and identify causative organisms. Therapy may be instituted before the results of the cultures and other laboratory studies are known; however, once these results become available, anti-infective therapy should be adjusted accordingly.

The species most frequently involved in causing human infections include *C. albicans*, *C. parapsilosis*, *C. tropicalis*, *C. glabrata* and *C. krusei*, all of which usually exhibit minimal inhibitory concentration (MICs) of less than 1 mg/L for voriconazole.

However, the *in vitro* activity of voriconazole against *Candida* species is not uniform. Specifically, for *C. glabrata*, the MICs of voriconazole for fluconazole-resistant isolates are proportionally higher than are those of fluconazole-susceptible isolates. Therefore, every attempt should be made to identify *Candida* to species level. If antifungal susceptibility testing is available, the MIC results may be interpreted using breakpoint criteria established by European Committee on Antimicrobial Susceptibility Testing (EUCAST).

EUCAST Breakpoints

Candida and Aspergillus species	Minimal Inhibitory Concentration (MIC) breakpoint (mg/L)	
	≤S (Susceptible)	>R (Resistant)
<i>Candida albicans</i> ¹	0.06	0.25
<i>Candida dubliniensis</i> ¹	0.06	0.25
<i>Candida glabrata</i>	Insufficient evidence (IE)	IE
<i>Candida krusei</i>	IE	IE
<i>Candida parapsilosis</i> ¹	0.125	0.25
<i>Candida tropicalis</i> ¹	0.125	0.25
<i>Candida guilliermondii</i> ²	IE	IE
Non-species related breakpoints for <i>Candida</i> ³	IE	IE
<i>Aspergillus fumigatus</i> ⁴	1	1
<i>Aspergillus nidulans</i> ⁴	1	1
<i>Aspergillus flavus</i>	IE ⁵	IE ⁵
<i>Aspergillus niger</i>	IE ⁵	IE ⁵

<i>Aspergillus terreus</i>	IE ⁵	IE ⁵
Non-species related breakpoints ⁶	IE	IE

¹ Strains with MIC values above the Susceptible/Intermediate (S/I) breakpoint are rare, or not yet reported. The identification and antifungal susceptibility tests on any such isolate must be repeated and if the result is confirmed the isolate sent to a reference laboratory. Until there is evidence regarding clinical response for confirmed isolates with MIC above the current resistant breakpoint they should be reported resistant. A clinical response of 76% was achieved in infections caused by the species listed below when MICs were lower than or equal to the epidemiological cut-offs. Therefore, wild type populations of *C. albicans*, *C. dubliniensis*, *C. parapsilosis* and *C. tropicalis* are considered susceptible.

² The epidemiological cut-off values (ECOFFs) for these species are in general higher than for *C. albicans*.

³ Non-species related breakpoints have been determined mainly on the basis of PK/PD data and are independent of MIC distributions of specific *Candida* species. They are for use only for organisms that do not have specific breakpoints.

⁴ Area of technical uncertainty (ATU) is 2. Report as R with the following comment: "In some clinical situations (non-invasive infections forms) voriconazole can be used provided sufficient exposure is ensured".

⁵ The ECOFFs for these species are in general one two-fold dilution higher than for *A. fumigatus*.

⁶ Non-species related breakpoints have not been determined.

Clinical experience

Successful outcome in this section is defined as complete or partial response.

Aspergillus infections – efficacy in aspergillosis patients with poor prognosis

Voriconazole has *in vitro* fungicidal activity against *Aspergillus* spp. The efficacy and survival benefit of voriconazole versus conventional amphotericin B in the primary treatment of acute invasive aspergillosis was demonstrated in an open, randomised, multicentre study in 277 immunocompromised patients treated for 12 weeks. Voriconazole was administered intravenously with a loading dose of 6 mg/kg every 12 hours for the first 24 hours followed by a maintenance dose of 4 mg/kg every 12 hours for a minimum of 7 days. Therapy could then be switched to the oral formulation at a dose of 200 mg every 12 hours. Median duration of IV voriconazole therapy was 10 days (range 2-85 days). After IV voriconazole therapy, the median duration of oral voriconazole therapy was 76 days (range 2-232 days).

A satisfactory global response (complete or partial resolution of all attributable symptoms, signs, radiographic/bronchoscopic abnormalities present at baseline) was seen in 53% of voriconazole-treated patients compared to 31% of patients treated with comparator. The 84-day survival rate for voriconazole was statistically significantly higher than that for the comparator and a clinically and statistically significant benefit was shown in favour of voriconazole for both time to death and time to discontinuation due to toxicity.

This study confirmed findings from an earlier, prospectively designed study where there was a positive outcome in subjects with risk factors for a poor prognosis, including graft versus host disease, and, in particular, cerebral infections (normally associated with almost 100% mortality).

The studies included cerebral, sinus, pulmonary and disseminated aspergillosis in patients with bone marrow and solid organ transplants, haematological malignancies, cancer and AIDS.

Candidaemia in non-neutropenic patients

The efficacy of voriconazole compared to the regimen of amphotericin B followed by fluconazole in the primary treatment of candidaemia was demonstrated in an open, comparative study. Three hundred and seventy non-neutropenic patients (above 12 years of age) with documented candidaemia were included in the study, of whom 248 were treated with voriconazole. Nine subjects in the voriconazole group and 5 in the amphotericin B followed by fluconazole group also had mycologically proven infection in deep tissue. Patients with renal failure were excluded from this study. The median treatment duration was 15 days in both treatment arms. In the primary analysis, successful response as assessed by a Data Review Committee (DRC) blinded to study medicinal product was defined as resolution/improvement in all clinical signs and symptoms of infection with eradication of *Candida* from blood and infected deep tissue sites 12 weeks after the end of therapy (EOT). Patients who did not have an assessment 12 weeks after EOT were counted as failures. In this analysis a successful response was seen in 41% of patients in both treatment arms.

In a secondary analysis, which utilised DRC assessments at the latest evaluable time point (EOT, or 2, 6, or 12 weeks after EOT) voriconazole and the regimen of amphotericin B followed by fluconazole had successful response rates of 65% and 71%, respectively.

The Investigator's assessment of successful outcome at each of these time points is shown in the following table.

<i>Timepoint</i>	<i>Voriconazole</i> (N=248)	<i>Amphotericin B fluconazole</i> (N=122)
EOT	178 (72%)	88 (72%)
2 weeks after EOT	125 (50%)	62 (51%)
6 weeks after EOT	104 (42%)	55 (45%)
12 weeks after EOT	104 (42%)	51 (42%)

Serious refractory *Candida* infections

The study comprised 55 patients with serious refractory systemic *Candida* infections (including candidaemia, disseminated and other invasive candidiasis) where prior antifungal treatment, particularly with fluconazole, had been ineffective. Successful response was seen in 24 patients (15 complete, 9 partial responses). In fluconazole-resistant non-*albicans* species, a successful outcome was seen in 3/3 *C. krusei* (complete responses) and 6/8 *C. glabrata* (5 complete, 1 partial response) infections. The clinical efficacy data were supported by limited susceptibility data.

Scedosporium and *Fusarium* infections

Voriconazole was shown to be effective against the following rare fungal pathogens:

Scedosporium spp.: Successful response to voriconazole therapy was seen in 16 (6 complete, 10 partial responses) of 28 patients with *S. apiospermum* and in 2 (both partial responses) of 7 patients with *S. prolificans* infection. In addition, a successful response was seen in 1 of 3 patients with infections caused by more than one organism including *Scedosporium* spp.

Fusarium spp.: Seven (3 complete, 4 partial responses) of 17 patients were successfully treated with voriconazole. Of these 7 patients, 3 had eye, 1 had sinus, and 3 had disseminated infection. Four additional patients with fusariosis had an infection caused by several organisms; 2 of them had a successful outcome.

The majority of patients receiving voriconazole treatment of the above-mentioned rare infections were intolerant of, or refractory to, prior antifungal therapy.

Paediatric population

Fifty-three paediatric patients aged 2 to <18 years were treated with voriconazole in two prospective, open-label, non-comparative, multi-center clinical trials. One study enrolled 31 patients with possible, proven or probable invasive aspergillosis (IA), of whom 14 patients had proven or probable IA and were included in the MITT efficacy analyses. The second study enrolled 22 patients with invasive candidiasis including candidaemia (ICC), and esophageal candidiasis (EC) requiring either primary or salvage therapy, of whom 17 were included in the MITT efficacy analyses. For patients with IA the overall rates of global response at 6 weeks were 64.3% (9/14), the global response rate was 40% (2/5) for patients 2 to <12 years and 77.8% (7/9) for patients 12 to <18 years of age. For patients with ICC the global response rate at EOT was 85.7% (6/7) and for patients with EC the global response rate at EOT was 70% (7/10). The overall rate of response (ICC and EC combined) was 88.9% (8/9) for 2 to <12 years old and 62.5% (5/8) for 12 to <18 years old.

Clinical studies examining QTc interval

A placebo-controlled, randomized, single-dose, crossover study to evaluate the effect on the QTc interval of healthy volunteers was conducted with three oral doses of voriconazole and ketoconazole. The placebo-adjusted mean maximum increases in QTc from baseline after 800, 1200 and 1600 mg of voriconazole were 5.1, 4.8, and 8.2 msec, respectively and 7.0 msec for ketoconazole 800 mg. No subject in any group had an increase in QTc of ≥ 60 msec from baseline. No subject experienced an interval exceeding the potentially clinically relevant threshold of 500 msec.

5.3 Pharmacokinetic properties

General pharmacokinetic characteristics

The pharmacokinetics of voriconazole have been characterised in healthy subjects, special populations and patients. During oral administration of 200 mg or 300 mg twice daily for 14 days in patients at risk of aspergillosis (mainly patients with malignant neoplasms of lymphatic or haematopoietic tissue), the observed pharmacokinetic characteristics of rapid and consistent absorption, accumulation and non-linear pharmacokinetics were in agreement with those observed in healthy subjects.

The pharmacokinetics of voriconazole are non-linear due to saturation of its metabolism. Greater than proportional increase in exposure is observed with increasing dose. It is estimated that, on average, increasing the oral dose from 200 mg twice daily to 300 mg twice daily leads to a 2.5-fold increase in exposure (AUC_{τ}). The oral maintenance dose of 200 mg (or 100 mg for patients

less than 40 kg) achieves a voriconazole exposure similar to 3 mg/kg IV. A 300 mg (or 150 mg for patients less than 40 kg) oral maintenance dose achieves an exposure similar to 4 mg/kg IV. When the recommended intravenous or oral loading dose regimens are administered, plasma concentrations close to steady-state are achieved within the first 24 hours of dosing. Without the loading dose, accumulation occurs during twice daily multiple dosing with steady-state plasma voriconazole concentrations being achieved by Day 6 in the majority of subjects.

Absorption

Voriconazole is rapidly and almost completely absorbed following oral administration, with maximum plasma concentrations (C_{max}) achieved 1-2 hours after dosing. The absolute bioavailability of voriconazole after oral administration is estimated to be 96%. When multiple doses of voriconazole are administered with high fat meals, C_{max} and AUC_{τ} are reduced by 34% and 24%, respectively. The absorption of voriconazole is not affected by changes in gastric pH.

Distribution

The volume of distribution at steady-state for voriconazole is estimated to be 4.6 L/kg, suggesting extensive distribution into tissues. Plasma protein binding is estimated to be 58%. Cerebrospinal fluid samples from eight patients in a compassionate programme showed detectable voriconazole concentrations in all patients.

Biotransformation

In vitro studies showed that voriconazole is metabolised by the hepatic cytochrome P450 isoenzymes, CYP2C19, CYP2C9 and CYP3A4.

The inter-individual variability of voriconazole pharmacokinetics is high.

In vivo studies indicated that CYP2C19 is significantly involved in the metabolism of voriconazole. This enzyme exhibits genetic polymorphism. For example, 15-20% of Asian populations may be expected to be poor metabolisers. For Caucasians and Blacks, the prevalence of poor metabolisers is 3-5%. Studies conducted in Caucasian and Japanese healthy subjects have shown that poor metabolisers have, on average, 4-fold higher voriconazole exposure (AUC_{τ}) than their homozygous extensive metaboliser counterparts. Subjects who are heterozygous extensive metabolisers have on average 2-fold higher voriconazole exposure than their homozygous extensive metaboliser counterparts.

The major metabolite of voriconazole is the N-oxide, which accounts for 72% of the circulating radiolabelled metabolites in plasma. This metabolite has minimal antifungal activity and does not contribute to the overall efficacy of voriconazole.

Elimination

Voriconazole is eliminated via hepatic metabolism with less than 2% of the dose excreted unchanged in the urine.

After administration of a radiolabelled dose of voriconazole, approximately 80% of the radioactivity is recovered in the urine after multiple intravenous dosing and 83% in the urine after

multiple oral dosing. The majority (>94%) of the total radioactivity is excreted in the first 96 hours after both oral and intravenous dosing.

The terminal half-life of voriconazole depends on dose and is approximately 6 hours at 200 mg (orally). Because of non-linear pharmacokinetics, the terminal half-life is not useful in the prediction of the accumulation or elimination of voriconazole.

Pharmacokinetics in special patient groups

Gender

In an oral multiple dose study, C_{max} and AUC_{τ} for healthy young females were 83% and 113% higher, respectively, than in healthy young males (18-45 years). In the same study, no significant differences in C_{max} and AUC_{τ} were observed between healthy elderly males and healthy elderly females (≥ 65 years).

In the clinical programme, no dosage adjustment was made on the basis of gender. The safety profile and plasma concentrations observed in male and female patients were similar. Therefore, no dosage adjustment based on gender is necessary.

Elderly

In an oral multiple dose study C_{max} and AUC_{τ} in healthy elderly males (≥ 65 years) were 61% and 86% higher, respectively, than in healthy young males (18-45 years). No significant differences in C_{max} and AUC_{τ} were observed between healthy elderly females (≥ 65 years) and healthy young females (18-45 years).

In the therapeutic studies no dosage adjustment was made on the basis of age. A relationship between plasma concentrations and age was observed. The safety profile of voriconazole in young and elderly patients was similar and, therefore, no dosage adjustment is necessary for the elderly (see section 4.2).

Paediatric population

The recommended doses in children and adolescent patients are based on a population pharmacokinetic analysis of data obtained from 112 immunocompromised paediatric patients aged 2 to <12 years and 26 immunocompromised adolescent patients aged 12 to <17 years. Multiple intravenous doses of 3, 4, 6, 7 and 8 mg/kg twice daily and multiple oral doses (using the powder for oral suspension) of 4 mg/kg, 6 mg/kg, and 200 mg twice daily were evaluated in 3 paediatric pharmacokinetic studies. Intravenous loading doses of 6 mg/kg IV twice daily on day 1 followed by 4 mg/kg intravenous dose twice daily and 300 mg oral tablets twice daily were evaluated in one adolescent pharmacokinetic study. Larger inter-subject variability was observed in paediatric patients compared to adults.

A comparison of the paediatric and adult population pharmacokinetic data indicated that the predicted total exposure (AUC_{τ}) in children following administration of a 9 mg/kg IV loading dose was comparable to that in adults following a 6 mg/kg IV loading dose.

The predicted total exposures in children following IV maintenance doses of 4 and 8 mg/kg twice daily were comparable to those in adults following 3 and 4 mg/kg IV twice daily, respectively. The predicted total exposure in children following an oral maintenance dose of 9 mg/kg (maximum of 350 mg) twice daily was comparable to that in adults following 200 mg oral twice daily. An 8 mg/kg intravenous dose will provide voriconazole exposure approximately 2-fold higher than a 9 mg/kg oral dose.

The higher intravenous maintenance dose in paediatric patients relative to adults reflects the higher elimination capacity in paediatric patients due to a greater liver mass to body mass ratio. Oral bioavailability may however be limited in paediatric patients with malabsorption and very low body weight for their age. In that case, intravenous voriconazole administration is recommended.

Voriconazole exposures in the majority of adolescent patients were comparable to those in adults receiving the same dosing regimens. However, lower voriconazole exposure was observed in some young adolescents with low body weight compared to adults. It is likely that these subjects may metabolize voriconazole more similarly to children than to adolescents/adults. Based on the population pharmacokinetic analysis, 12 to 14-year-old adolescents weighing less than 50 kg should receive children's doses (see section 4.2).

Renal impairment

Tablets:

In an oral single-dose (200 mg) study in subjects with normal renal function and mild (creatinine clearance 41-60 ml/min) to severe (creatinine clearance <20 ml/min) renal impairment, the pharmacokinetics of voriconazole were not significantly affected by renal impairment. The plasma protein binding of voriconazole was similar in subjects with different degrees of renal impairment (see sections 4.2 and 4.4).

Powder for solution for infusion:

In patients with moderate to severe renal dysfunction (serum creatinine levels >2.5 mg/dl), accumulation of the intravenous vehicle, SBECD, occurs (see sections 4.2 and 4.4).

Hepatic impairment

After an oral single-dose (200 mg), AUC was 233% higher in subjects with mild to moderate hepatic cirrhosis (Child-Pugh A and B) compared with subjects with normal hepatic function. Protein binding of voriconazole was not affected by impaired hepatic function.

In an oral multiple-dose study, AUC_τ was similar in subjects with moderate hepatic cirrhosis (Child-Pugh B) given a maintenance dose of 100 mg twice daily and subjects with normal hepatic function given 200 mg twice daily. No pharmacokinetic data are available for patients with severe hepatic cirrhosis (Child-Pugh C) (see sections 4.2 and 4.4).

6. NON-CLINICAL PROPERTIES

6.1. Animal toxicology or pharmacology

Repeated-dose toxicity studies with voriconazole indicated the liver to be the target organ. Hepatotoxicity occurred at plasma exposures similar to those obtained at therapeutic doses in humans, in common with other antifungal agents. In rats, mice and dogs, voriconazole also induced minimal adrenal changes. Conventional studies of safety pharmacology, genotoxicity or carcinogenic potential did not reveal a special hazard for humans.

In reproduction studies, voriconazole was shown to be teratogenic in rats and embryotoxic in rabbits at systemic exposures equal to those obtained in humans with therapeutic doses. In the pre- and post-natal development study in rats at exposures lower than those obtained in humans with therapeutic doses, voriconazole prolonged the duration of gestation and labour and produced dystocia with consequent maternal mortality and reduced perinatal survival of pups. The effects on parturition are probably mediated by species-specific mechanisms, involving reduction of oestradiol levels, and are consistent with those observed with other azole antifungal agents. Voriconazole administration induced no impairment of male or female fertility in rats at exposures similar to those obtained in humans at therapeutic doses.

Preclinical data on the intravenous vehicle SBECD indicated that the main effects were vacuolation of urinary tract epithelium and activation of macrophages in the liver and lungs in the repeated-dose toxicity studies. As GPMT (guinea pig maximisation test) result was positive, prescribers should be aware of the hypersensitivity potential of the intravenous formulation. Standard genotoxicity and reproduction studies with the excipient SBECD reveal no special hazard for humans. Carcinogenicity studies were not performed with SBECD. An impurity present in SBECD has been shown to be an alkylating mutagenic agent with evidence for carcinogenicity in rodents. This impurity should be considered a substance with carcinogenic potential in humans. In light of these data the duration of treatment with the intravenous formulation should be no longer than 6 months.

7. DESCRIPTION

Tablets:

Voriconazole 50 mg film-coated tablets are white, round tablets, debossed "Pfizer" on one side and "VOR50" on the reverse.

Voriconazole 200 mg film-coated tablets are white, capsule-shaped tablets, debossed "Pfizer" on one side and "VOR200" on the reverse.

Powder for solution for infusion:

Voriconazole powder for solution for infusion is a white lyophilised powder containing nominally 200 mg voriconazole presented in a 30 ml clear glass vial.

8. PHARMACEUTICAL PARTICULARS

8.1 Incompatibilities

Film-coated tablets:

Not applicable

Powder for solution for infusion:

Voriconazole must not be infused into the same line or cannula concomitantly with other intravenous products. The bag should be checked to ensure that the infusion is complete. When the voriconazole infusion is complete, the line may be used for administration of other intravenous products.

Blood products and concentrated electrolytes

Voriconazole must not be infused concomitantly with any blood product or any short-term infusion of concentrated electrolytes, even if the two infusions are running in separate intravenous lines (or cannulas). Electrolyte disturbances such as hypokalemia, hypomagnesemia and hypocalcemia should be corrected prior to initiation of voriconazole therapy (see sections 4.2 and 4.4).

Intravenous solutions containing (non-concentrated) electrolytes

Voriconazole can be infused at the same time as other intravenous solutions containing (non-concentrated) electrolytes, but must be infused through a separate line.

Total parenteral nutrition (TPN)

Voriconazole can be infused at the same time as total parenteral nutrition, but must be infused in a separate line. If infused through a multiple-lumen catheter, TPN needs to be administered using a different port from the one used for voriconazole.

Voriconazole must not be diluted with 4.2% Sodium Bicarbonate Infusion. Compatibility with other concentrations is unknown.

This medicinal product must not be mixed with other medicinal products except those mentioned in section 8.4.

8.2 Shelf-life

Film-coated tablets:

3 years

Powder for solution for infusion:

3 years

8.3 Packaging information

Film-coated tablets:

Monocarton containing a blister pack of 7 tablets.

Blister pack:

50 mg tablets: Blister pack made up of 250 microns transparent PVC and 25 microns hard tempered aluminium with vinyl-acrylate based heat seal coating on the bright side of the foil, 1.6 g/m² polyester based lacquer on the matt side.

200 mg tablets: Blister pack made up of 300 microns transparent PVC and 25 microns hard tempered aluminium with vinyl-acrylate based heat seal coating on the bright side of the foil, 1.6 g/m² polyester based lacquer on the matt side.

Powder for solution for infusion:

Voriconazole is supplied as a sterile lyophilised powder in individually boxed, single use 30 ml clear Type I glass vials with rubber stoppers and aluminium caps with plastic seals.

8.4. Storage and handling instructions

Film-coated tablets:

Store in a dry, well-ventilated place at a temperature not exceeding 30°C.
Keep out of reach of children.

Powder for solution for infusion:

The unreconstituted vial should be stored in a dry, well-ventilated place at a temperature not exceeding 30°C.

Reconstituted Concentrate: Store at 2°C-8°C for up to 24 hours (in a refrigerator).

Instructions for Use and Handling

Film-coated tablets:

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Powder for solution for infusion:

Voriconazole is supplied in single use vials. The vial contents are reconstituted with 19 ml of Water for Injections to obtain a clear solution containing 10 mg/ml of voriconazole and an extractable volume of 20 ml. Discard the vial if vacuum does not pull the diluent into the vial. For administration, the required volume of the reconstituted solution (table below) is added to a recommended compatible infusion solution (detailed below) to obtain, where appropriate, a final voriconazole solution containing 0.5-5 mg/ml.

The reconstituted solution can be diluted with:

Sodium Chloride 9 mg/ml (0.9%) Solution for Injection
Compound Sodium Lactate Intravenous Infusion
5% Glucose and Lactated Ringer's Intravenous Infusion
5% Glucose and 0.45% Sodium Chloride Intravenous Infusion
5% Glucose Intravenous Infusion
5% Glucose in 20 mEq Potassium Chloride Intravenous Infusion
0.45% Sodium Chloride Intravenous Infusion
5% Glucose and 0.9% Sodium Chloride Intravenous Infusion

The compatibility of voriconazole with diluents other than described above or in section 8.1 is unknown.

Required Volumes of 10 mg/ml Voriconazole Concentrate:

Body Weight (kg)	Volume of voriconazole Concentrate (10 mg/ml) required for:				
	3 mg/kg dose (number of vials)	4 mg/kg dose (number of vials)	6 mg/kg dose (number of vials)	8 mg/kg dose (number of vials)	9 mg/kg dose (number of vials)
10	-	4.0 ml (1)	-	8.0 ml (1)	9.0 ml (1)
15	-	6.0 ml (1)	-	12.0 ml (1)	13.5 ml (1)
20	-	8.0 ml (1)	-	16.0 ml (1)	18.0 ml (1)
25	-	10.0 ml (1)	-	20.0 ml (1)	22.5 ml (2)
30	9.0 ml (1)	12 ml (1)	18 ml (1)	24.0 ml (2)	27.0 ml (2)
35	10.5 ml (1)	14 ml (1)	21 ml (2)	28.0 ml (2)	31.5 ml (2)
40	12.0 ml (1)	16 ml (1)	24 ml (2)	32.0 ml (2)	36.0 ml (2)
45	13.5 ml (1)	18 ml (1)	27 ml (2)	36.0 ml (2)	40.5 ml (3)
50	15.0 ml (1)	20 ml (1)	30 ml (2)	40.0 ml (2)	45.0 ml (3)
55	16.5 ml (1)	22 ml (2)	33 ml (2)	44.0 ml (3)	49.5 ml (3)
60	18.0 ml (1)	24 ml (2)	36 ml (2)	48.0 ml (3)	54.0 ml (3)
65	19.5 ml (1)	26 ml (2)	39 ml (2)	52.0 ml (3)	58.5 ml (3)
70	21.0 ml (2)	28 ml (2)	42 ml (3)	-	-
75	22.5 ml (2)	30 ml (2)	45 ml (3)	-	-
80	24.0 ml (2)	32 ml (2)	48 ml (3)	-	-
85	25.5 ml (2)	34 ml (2)	51 ml (3)	-	-
90	27.0 ml (2)	36 ml (2)	54 ml (3)	-	-
95	28.5 ml (2)	38 ml (2)	57 ml (3)	-	-
100	30.0 ml (2)	40 ml (2)	60 ml (3)	-	-

Voriconazole is a single dose unpreserved sterile lyophile. Therefore, from a microbiological point of view, the product must be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C to 8°C, unless reconstitution has taken place in controlled and validated aseptic conditions.

Discard the reconstituted solution if content is not clear or if particulate matter or discoloration is observed.

9. PATIENT COUNSELLING INFORMATION

Visual Disturbances

Patients should be instructed that visual disturbances such as blurring and sensitivity to light may occur with the use of VFEND.

Photosensitivity

- Advise patients of the risk of photosensitivity (with or without concomitant methotrexate), accelerated photoaging, and skin cancer.

- Advise patients that VFEND can cause serious photosensitivity and to immediately contact their healthcare provider for new or worsening skin rash.
- Advise patients to avoid exposure to direct sun light and to use measures such as protective clothing and sunscreen with high sun protection factor (SPF).

Embryo-Fetal Toxicity

- Advise female patients of the potential risks to a fetus.
- Advise females of reproductive potential to use effective contraception during treatment with Voriconazole.

10. DETAILS OF MANUFACTURER

Please refer to outer carton

11. DETAILS OF PERMISSION OR LICENCE NUMBER WITH DATE

1732/04 dated 8-Sep-2004

12. DATE OF REVISION

March 2026