

*Your doctor is the best resource for medical advice and information. The health information contained herein is provided for educational / awareness purposes only and is not intended to replace discussions with a medical practitioner and/or medical advice or be construed as a promotional information.*

# Package Leaflet: Information for the user

## Sunitinib Malate Capsules

### Sutent<sup>®</sup>

---

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- If you have any further questions, ask your doctor.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Sutent (Sunitinib Malate) is and what it is used for
2. What you need to know before you take Sutent (Sunitinib Malate)
3. How to take Sutent (Sunitinib Malate)
4. Possible side effects
5. How to store Sutent (Sunitinib Malate)
6. Contents of the pack and other information

#### **1. What Sutent (Sunitinib Malate) is and what it is used for**

Sutent (Sunitinib Malate) contains the active substance sunitinib, which is a protein kinase inhibitor. It is used to treat cancer by preventing the activity of a special group of proteins which are known to be involved in the growth and spread of cancer cells.

Sutent (Sunitinib Malate) is used to treat patients with the following types of cancer:

- Gastrointestinal stromal tumour (GIST), a type of cancer of the stomach and bowel, where imatinib mesylate (another anticancer medicine) no longer works or you cannot take imatinib.
- Advanced renal cell carcinoma, a type of kidney cancer that has spread to other parts of the body.

---

® Trademark Proprietor: C. P. Pharmaceuticals International C.V.  
Licensed User: Pfizer Products India Private Limited, India

- Metastatic well-differentiated pancreatic neuroendocrine tumours (tumours of the hormone-producing cells in the pancreas) that have progressed in adults or cannot be removed with surgery.

If you have any questions about how Sutent (Sunitinib Malate) works or why this medicine has been prescribed for you, ask your doctor.

## 2. What you need to know before you take Sutent (Sunitinib Malate)

### Do not take Sutent (Sunitinib Malate)

- If you are allergic to sunitinib or any of the other ingredients of Sutent (Sunitinib Malate) (listed in section 6).

### Warnings and precautions

#### Talk to your doctor before taking Sutent (Sunitinib Malate):

- **If you have high blood pressure.** Sutent (Sunitinib Malate) can raise blood pressure. Your doctor may check your blood pressure during treatment with Sutent (Sunitinib Malate), and you may be treated with medicines to reduce the blood pressure, if needed.
- **If you have or have had blood disease, bleeding problems, or bruising.** Treatment with Sutent (Sunitinib Malate) may lead to a higher risk of bleeding and/or gastrointestinal bleeding, respiratory bleeding, urinary tract bleeding, cell death and bleeding in the brain; and may lead to changes in the number of certain cells in the blood which may lead to anaemia or decrease neutrophil and platelet count, affect the ability of your blood to clot. If you are taking warfarin or acenocoumarole, medicines which thin the blood to prevent blood clots, there may be a greater risk of bleeding. Tell your doctor if you have any bleeding while on treatment with Sutent (Sunitinib Malate).
- **If you have heart problems.** Sutent (Sunitinib Malate) can cause heart problems. Tell your doctor if you have shortness of breath, or have swollen feet and ankles. It may lead to a disease of the heart muscle that makes it difficult to pump blood (cardiomyopathy).
- **If you have abnormal heart rhythm changes.** Sutent (Sunitinib Malate) can cause abnormality of your heart rhythm and QT prolongation on ECG. Your doctor may obtain electrocardiograms to evaluate for these problems during your treatment with Sutent (Sunitinib Malate). Tell your doctor if you feel dizzy, faint, or have abnormal heartbeats while taking Sutent (Sunitinib Malate).
- **If you have had a recent problem with blood clots in your veins and/or arteries (types of blood vessels), including stroke, heart attack, embolism, or thrombosis.** Call your doctor immediately if you get symptoms such as chest pain or pressure, pain in your arms, back, neck or jaw, shortness of breath, numbness or weakness on 1 side of your body, trouble talking, headache, or dizziness while on treatment with Sutent (Sunitinib Malate).

- **If you have or have had an aneurysm** (enlargement and weakening of a blood vessel wall) or a tear in a blood vessel wall.
- **If you have or have had damage to the smallest blood vessels known as thrombotic microangiopathy (TMA).**
- **If you have thyroid glands problems.** Sutent (Sunitinib Malate) can cause thyroid gland problems. Tell your doctor if you get tired more easily, generally feel colder than other people, or your voice deepens, you gain or loss weight, increase appetite, sweating whilst taking Sutent (Sunitinib Malate). Your thyroid function should be checked before you take Sutent (Sunitinib Malate) and regularly while you are taking it. If your thyroid gland is not producing enough thyroid hormone, you may be treated with thyroid hormone replacement.
- **If you have or have had pancreatic disorders.** Tell your doctor if you develop any of the following signs and symptoms: pain in the area of the stomach (upper abdomen) or back, nausea, vomiting, and fever. These may be caused by inflammation of the pancreas or gallbladder.
- **If you have or have had liver problems.** Tell your doctor if you develop any of the following signs and symptoms of liver problems during Sutent (Sunitinib Malate) treatment: itching, yellow eyes or skin, dark urine, pale stool and pain or discomfort in the right upper stomach area, fever, nausea. Your doctor should do blood tests to check your liver function before and during treatment with Sutent (Sunitinib Malate), and as clinically indicated.
- **If you have or have had kidney problems.** Your doctor will monitor your kidney function. Baseline urinalysis is recommended, and patients should be monitored for the development or worsening of proteinuria.
- **If you are going to have surgery or if you had an operation recently.** Sutent (Sunitinib Malate) may affect the way your wounds heal. You will usually be taken off Sutent (Sunitinib Malate) if you are having an operation. Your doctor will decide when to start Sutent (Sunitinib Malate) again.
- **You may be advised to have a dental check-up before you start treatment with Sutent (Sunitinib Malate).**
  - If you have or have had osteonecrosis of jaw, pain in the mouth, teeth and/or jaw, swelling or sores inside the mouth, numbness or a feeling of heaviness in the jaw, or loosening of a tooth, tell your doctor and dentist immediately.
  - If you need to undergo an invasive dental treatment or dental surgery, tell your dentist that you are being treated with Sutent (Sunitinib Malate) in particular when you are also receiving or have received intravenous bisphosphonates. Bisphosphonates are

medicines used to prevent bone complications that may have been given for another medical condition.

- **If you have or have had skin and subcutaneous tissue disorders.** While you are on this medicine "pyoderma gangrenosum" (painful skin ulceration) or "necrotising fasciitis" (rapidly spreading infection of the skin/soft tissue that may be life-threatening) may occur. Contact your doctor immediately if symptoms of infection occur around a skin injury, including fever, pain, redness, swelling, or drainage of pus or blood. This event is generally reversible after sunitinib discontinuation. Severe skin rashes (Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme) have been reported with the use of sunitinib, appearing initially as reddish target-like spots or circular patches often with central blisters on the trunk. The rash may progress to widespread blistering or peeling of the skin and may be life-threatening. If you develop a rash or these skin symptoms, seek immediate advice from a doctor. Patients should be advised that depigmentation of the hair or skin may occur during treatment with sunitinib. Other possible dermatological effects may include dryness, thickness or cracking of the skin, blisters, or rash on the palms of the hands and soles of the feet.
- **If you have or have had seizures.** Notify your doctor as soon as possible if you have high blood pressure, headache, or loss of sight, decreased alertness, altered mental functioning.
- **If you have diabetes.** Blood sugar levels in diabetic patients should be checked regularly in order to assess if antidiabetic medicine's dosage needs to be adjusted to minimise the risk of low blood sugar. Notify your doctor as soon as possible if you experience any signs and symptoms of low blood sugar (fatigue, palpitations, sweating, hunger and loss of consciousness).
- **If you have had Tumor lysis syndrome (TLS).** A condition that occurs when a large number of cancer cells die within a short period, releasing their contents in to the blood; your doctor may consider stopping the medication.
- **If you have had Serious infections,** inform your doctor.
- **If fistula formation** (when inflammation causes sores, or ulcers, to form on the inside wall of the intestine or nearby organs) occurs, sunitinib treatment should be interrupted.
- **If undesirable reactions (hypersensitivity)** lead to rapid swelling, of the area beneath the skin or mucosa (angioedema - swelling of the face and throat), notify your doctor at the earliest.

### Children and adolescents

Sutent (Sunitinib Malate) is not recommended for people aged under 18.

### Other medicines and Sutent (Sunitinib Malate)

Tell your doctor if you are taking, have recently taken, or might take any other medicines, including medicines obtained without a prescription and even those not prescribed.

Some medicines can affect the levels of Sutent (Sunitinib Malate) in your body. You should inform your doctor if you are taking medicines containing the following active substances:

- ketoconazole, itraconazole
- erythromycin, clarithromycin, rifampicin
- ritonavir
- dexamethasone – a corticosteroid used for various conditions
- phenytoin, carbamazepine, phenobarbital
- herbal preparations containing St. John’s Wort (*Hypericum perforatum*)

### **Sutent (Sunitinib Malate) with food and drink**

You should avoid drinking grapefruit juice while on treatment with Sutent (Sunitinib Malate).

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant, or are planning to have a baby, ask your doctor for advice before taking this medicine.

If you might get pregnant, you should use a reliable method of contraception during treatment with Sutent (Sunitinib Malate).

If you are breast-feeding, tell your doctor. You should not breast-feed during treatment with Sutent (Sunitinib Malate).

### **Fertility**

Male and female fertility may be compromised by treatment with sunitinib.

### **Driving and using machines**

If you experience dizziness or you feel unusually tired, take special care when driving or using machines.

### **Sutent (Sunitinib Malate) contains sodium**

This medicine contains less than 1 mmol (23 mg) sodium per capsule, that is to say essentially ‘sodium-free’.

## **3. How to take Sutent (Sunitinib Malate)**

Always take this medicine exactly as your doctor has told you. Check with your doctor if you are not sure.

Your doctor will prescribe a dose that is right for you, depending on the type of cancer to be treated. If you are being treated for:

- GIST or MRCC: the usual dose is 50 mg once daily taken for 28 days (4 weeks), followed by 14 days (2 weeks) of rest (no medicine), in 6-week cycles.
- pNET: the usual dose is 37.5 mg once daily without a rest period.

Your doctor will determine the appropriate dose you need to take, as well as if and when you need to stop treatment with Sutent (Sunitinib Malate).

Sutant (Sunitinib Malate) can be taken with or without food.

### **If you take more Sutent (Sunitinib Malate) than you should**

If you have accidentally taken too many capsules, talk to your doctor straight away. You may require medical attention.

### **If you forget to take Sutent (Sunitinib Malate)**

Do not take a double dose to make up for a forgotten dose.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

You must immediately contact your doctor if you experience any of those serious side effects (see also **What you need to know before you take Sutent (Sunitinib Malate)**):

**Heart problems.** Tell your doctor if you feel very tired, are short of breath, or have swollen feet and ankles. These may be symptoms of heart problems that may include heart failure and heart muscle problems (cardiomyopathy).

**Lung or breathing problems.** Tell your doctor if you develop cough, chest pain, sudden onset of shortness of breath, or coughing up blood. These may be symptoms of a condition called pulmonary embolism that occurs when blood clots travel to your lungs.

**Kidney disorders.** Tell your doctor if you experience altered frequency or absence of urination which may be symptoms of kidney failure.

**Bleeding.** Tell your doctor if you have any of these symptoms or a serious bleeding problem during treatment with Sutent (Sunitinib Malate): painful, swollen stomach (abdomen); vomiting blood; black, sticky stools; bloody urine; headache or change in your mental status; coughing up of blood or bloody sputum from the lungs or airway.

**Tumour destruction leading to hole in the intestine.** Tell your doctor if you have severe abdominal pain, fever, nausea, vomiting, blood in your stool, or changes in bowel habits.

Other side effects with Sutent (Sunitinib Malate) may include:

Very common: may affect more than 1 in 10 people

- Reduction in the number of platelets, red blood cells and/or white blood cells (e.g. neutrophils).
- Shortness of breath.
- High blood pressure.
- Extreme tiredness, loss of strength.
- Swelling caused by fluid under the skin and around the eye, deep allergic rash.
- Mouth pain/irritation, mouth sores/inflammation/dryness, taste disturbances, upset stomach, nausea, vomiting, diarrhoea, constipation, abdominal pain/swelling, loss/decrease of appetite.
- Decreased activity of thyroid gland (hypothyroidism).
- Dizziness.
- Headache.
- Nose bleeding.
- Back pain, joint pain.
- Pain in arms and legs.
- Yellow skin/skin discoloration, excess pigmentation of the skin, hair colour change, rash on the palms of the hands and soles of the feet, rash, dryness of the skin, Mucosal inflammation.
- Cough.
- Fever.
- Difficulty in falling asleep.

Common: may affect up to 1 in 10 people

- Blood clots in the blood vessels.
- Deficiency of blood supply to the heart muscle, due to obstruction or constriction of the coronary arteries.
- Chest pain, shortness of breath (dyspnea), oropharyngeal pain.
- Decreased in the amount of blood pumped by the heart.
- Coughing up of blood (hemoptysis).
- Fluid retention including around the lungs.
- Infections.
- Complication of severe infection (infection is present in the bloodstream) that can lead to tissue damage, organ failure, and death.
- Decreased blood sugar level (see section 2).
- Loss of protein in the urine sometime resulting in swelling.
- Influenza-like syndrome.
- Abnormal blood tests including pancreatic and liver enzymes, kidney functions.
- High level of uric acid in the blood.
- Haemorrhoids, pain in the rectum, gingival bleeding, difficulty in swallowing or inability to swallow.
- Burning or painful sensation in the tongue, inflammation of the digestive tract lining, excessive gas in the stomach or intestine.
- Weight loss.

- Swelling around the eyes (Periorbital oedema), fluid retention around the eye (eyelid edema).
- Musculoskeletal pain (pain in muscles and bones), muscular weakness, muscular fatigue, muscle pain, muscle spasms.
- Nasal dryness, congested nose.
- Excessive tear flow.
- Abnormal sensation of the skin, itching, flaking and inflammation of the skin, blisters, acne, nail discolouration, hair loss.
- Abnormal sensations in extremities.
- Abnormally decreased/increased sensitivity, particularly to touch.
- Acid heartburn.
- Dehydration.
- Hot flushes.
- Abnormally coloured urine.
- Depression.
- Chills.

Uncommon: may affect up to 1 in 100 people

- Stroke.
- Heart attack caused by an interrupted or decreased blood supply to the heart.
- Changes in the electrical activity or abnormal rhythm of the heart.
- Fluid around the heart (pericardial effusion).
- Uncontrolled bleeding in the brain (cerebral hemorrhage), stopped blood flow to a part of the brain either by a blockage or the rupture of a blood vessel (cerebrovascular accident), a temporary period of symptoms similar to those of a stroke (Transient ischemic attack), QT prolongation on ECG, uncontrolled bleeding of the tumor (Tumor hemorrhage), acute bleeding from the lungs (pulmonary hemorrhage), acute bleeding in the urinary tract (urinary tract hemorrhage).
- Liver failure.
- Pain in the stomach (abdomen) caused by inflammation of the pancreas.
- Tumour destruction leading to hole in the intestine (perforation).
- Inflammation (swelling and redness) of the gallbladder with or without associated gallstones.
- Abnormal tube like passage from one normal body cavity to another body cavity or the skin.
- Pain in the mouth, teeth and/or jaw, swelling or sores inside the mouth, numbness or a feeling of heaviness in the jaw, or loosening of a tooth. These could be signs and symptoms of bone damage in the jaw (osteonecrosis), see section 2.
- Overproduction of thyroid hormones which increases the amount of energy the body uses at rest.
- Problems with wound healing after surgery.
- Increased blood level of enzyme (creatine phosphokinase) from muscle.
- Excessive reaction to an allergen including hay fever, skin rash, itchy skin, hives, swelling of body parts, and trouble breathing.
- Inflammation of the colon (colitis, colitis ischaemic).



Rare: may affect up to 1 in 1,000 people

- Severe reaction of the skin and/or mucous membranes (Stevens-Johnson syndrome, toxic epidermal necrolysis, erythema multiforme).
- Tumour lysis syndrome (TLS) – TLS consists of a group of metabolic complications that can occur during treatment of cancer. These complications are caused by the break-down products of dying cancer cells and may include the following: nausea, shortness of breath, irregular heartbeat, muscular cramps, seizure, clouding of urine and tiredness associated with abnormal laboratory test results (high potassium, uric acid and phosphorous levels and low calcium levels in the blood) that can lead to changes in kidney function and acute renal failure.
- Abnormal muscle breakdown which can lead to kidney problems (rhabdomyolysis).
- Reversible posterior leukoencephalopathy syndrome.
- Nephrotic syndrome
- Rapid edema, or swelling, of the area beneath the skin or mucosa (angioedema)
- Disease that affects the muscles that control voluntary movement in the body (myopathy)
- Left ventricular failure
- *Torsade de pointes*
- Painful skin ulceration (pyoderma gangrenosum).
- Inflammation of the liver (hepatitis).
- Inflammation of the thyroid gland.
- Damage to the smallest blood vessels known as thrombotic microangiopathy (TMA).

Not known (frequency cannot be estimated from available data):

- An enlargement and weakening of a blood vessel wall or a tear in a blood vessel wall (aneurysms and artery dissections).
- Lack of energy, confusion, sleepiness, unconsciousness/coma – these symptoms may be signs of brain toxicity caused by high blood levels of ammonia (hyperammonaemic encephalopathy).

##### **5. How to store Sutent (Sunitinib Malate)**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on carton, bottle and blister foil after “EXP”. The expiry date refers to the last day of that month.
- Store below 30°C in the original package. Protect from moisture.
- Do not use this medicine if you notice that the pack is damaged or shows signs of tampering.

Do not throw away any medicines via wastewater or household waste. Ask your doctor how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Sutent (Sunitinib Malate) contains

#### Sutent (Sunitinib Malate) 12.5 mg hard capsules

The active substance is sunitinib. Each capsule contains sunitinib malate equivalent to 12.5 mg sunitinib. The other ingredients are:

- *Capsule content:* Mannitol, Croscarmellose Sodium, Povidone (K-25), Magnesium stearate.
- *Capsule shell:* Gelatin; Red iron oxide (E172), Titanium dioxide (E171).
- *Printing ink:* Shellac, Propylene glycol, Sodium hydroxide, Povidone, Titanium dioxide (E171).

#### Sutent (Sunitinib Malate) 25 mg hard capsules

The active substance is sunitinib. Each capsule contains sunitinib malate equivalent to 25 mg sunitinib.

The other ingredients are:

- *Capsule content:* Mannitol (E421), Croscarmellose sodium, Povidone (K-25), Magnesium stearate.
- *Capsule shell:* Gelatin, Red iron oxide (E172), Titanium dioxide (E171), Yellow iron oxide (E172), Black iron oxide (E172).
- *Printing ink:* Shellac, Propylene glycol, Sodium hydroxide, Povidone, Titanium dioxide (E171).

#### Sutent (Sunitinib Malate) 50 mg hard capsules

The active substance is sunitinib. Each capsule contains sunitinib malate equivalent to 50 mg sunitinib.

The other ingredients are:

- *Capsule content:* Mannitol (E421), Croscarmellose sodium, Povidone (K-25), Magnesium stearate.
- *Capsule shell:* Gelatin, Titanium dioxide (E171), Yellow iron oxide (E172), Red iron oxide (E172), Black iron oxide (E172).
- *Printing ink:* Shellac, Propylene glycol, Sodium hydroxide, Povidone, Titanium dioxide (E171).

### What Sutent (Sunitinib Malate) looks like and contents of the pack

Sutent (Sunitinib Malate) 12.5 mg is supplied as hard gelatin capsules with orange cap and orange body, printed with white ink “Pfizer” on the cap, “STN 12.5 mg” on the body, and containing yellow to orange granules.

Sutent (Sunitinib Malate) 25 mg is supplied as hard gelatin capsules with caramel cap and orange body, printed with white ink “Pfizer” on the cap, “STN 25 mg” on the body, and containing yellow to orange granules.

Sutent (Sunitinib Malate) 50 mg is supplied as hard gelatin capsules with caramel cap and caramel body, printed with white ink “Pfizer” on the cap, “STN 50 mg” on the body, and containing yellow to orange granules.

It is available in bottles of 28 capsules and in blister pack of 7 capsules.

All strengths/pack sizes mentioned in this document might not be available in the market.

**Imported and marketed by**

Pfizer Products India Private Limited,  
The Capital- B Wing, 1802, 18<sup>th</sup> Floor,  
Plot No. C-70, G Block, Bandra Kurla Complex,  
Bandra (East), Mumbai 400 051,  
India

**Manufacturer**

M/s. Pfizer Italia S.r.l. Localita Marino del Tronto Ascoli Piceno - 63100 (Italy)

*This leaflet was prepared based on LPD no. LPDSUT042024*