

Alavert Allergy Sinus D12

Loratadine, Pseudoephedrine Sulfate

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Drug Facts	
Active ingredients (in each tablet)	Purposes
Loratadine 5 mg.....	Antihistamine
Pseudoephedrine sulfate 120 mg.....	Nasal decongestant

Uses
<ul style="list-style-type: none">▪ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:<ul style="list-style-type: none">▪ runny nose▪ sneezing▪ itchy, watery eyes▪ itching of the nose or throat▪ temporarily relieves nasal congestion due to the common cold, hay fever or other respiratory allergies▪ reduces swelling of nasal passages▪ temporarily relieves sinus congestion and pressure▪ temporarily restores freer breathing through the nose

Warnings
Do not use <ul style="list-style-type: none">▪ if you have ever had an allergic reaction to this product or any of its ingredients▪ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Alavert Allergy Sinus D12

Loratadine, Pseudoephedrine Sulfate

This is the most current labeling information and may differ from labels on product packaging. If there are any differences between this website labeling and product packaging labeling, this website labeling should be regarded as the most current.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not divide, crush, chew or dissolve the tablet

Age	Dose
adults and children 12 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- **each tablet contains:** calcium 30 mg
- store between 15° and 25°C (59° and 77°F)
- keep in a dry place

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

Questions or comments? Call weekdays from 9 AM to 5 PM EST at **1-800-ALAVERT (1-800-252-8378)**